

On the Inextendibility of Spacetime*

John Byron Manchak

[Forthcoming in *Philosophy of Science*.]

Abstract

It has been argued that spacetime must be inextendible – that it must be “as large as it can be” in some sense. Here, we register some skepticism with respect to this position.

1 Introduction

A spacetime is counted as inextendible if, intuitively, it is “as large as it can be”. It has been argued that inextendibility is a “reasonable physical condition to be imposed on models of the universe” (Geroch 1970, 262) and that a spacetime must be inextendible if it is “to be a serious candidate for describing actuality” (Earman 1995, 32). Here, in a variety of ways, we register some skepticism with respect to such positions.

2 Preliminaries

We begin with a few preliminaries concerning the relevant background formalism of general relativity.¹ An n -dimensional, relativistic *spacetime* (for $n \geq 2$) is a pair of mathematical objects (M, g_{ab}) where M is a connected n -dimensional Hausdorff manifold (without boundary) that is smooth and g_{ab} is a smooth, non-degenerate, pseudo-Riemannian metric of Lorentz signature $(-, +, \dots, +)$ defined on M . We say two spacetimes (M, g_{ab}) and (M', g'_{ab}) are *isometric* if there is a diffeomorphism $\varphi : M \rightarrow M'$ such that $\varphi_* g_{ab} = g'_{ab}$. Two spacetimes (M, g_{ab}) and (M', g'_{ab}) are *locally isometric* if, for each point $p \in M$, there is an open neighborhood

*Thanks to Thomas Barrett, Ben Feintzeig, David Malament, Jim Weatherall, and Chris Wüthrich for help with earlier versions of this document. Special thanks go to David Malament for providing a sketch of the Lemma 2 proof.

¹The reader is encouraged to consult Hawking and Ellis (1973), Wald (1984), and Malament (2012) for details. An outstanding (and less technical) survey of the global structure of spacetime is given by Geroch and Horowitz (1979).

O of p and an open subset O' of M' such that O and O' are isometric, and, correspondingly, with the roles of (M, g_{ab}) and (M', g'_{ab}) interchanged.

A spacetime (M, g_{ab}) is *extendible* if there exists a spacetime (M', g'_{ab}) and a proper isometric embedding $\varphi : M \rightarrow M'$. Here, the spacetime (M', g'_{ab}) is an *extension* of (M, g_{ab}) . A spacetime is *inextendible* if it has no extension. A \mathcal{P} -spacetime is a spacetime with property \mathcal{P} . A \mathcal{P} -spacetime (M', g'_{ab}) is a \mathcal{P} -*extension* of a \mathcal{P} -spacetime (M, g_{ab}) if (M', g'_{ab}) is an extension of (M, g_{ab}) . A \mathcal{P} -spacetime is \mathcal{P} -*extendible* if it has a \mathcal{P} -extension and is \mathcal{P} -*inextendible* otherwise. We say (M, g_{ab}, F) is an n -dimensional *framed spacetime* if (M, g_{ab}) is an n -dimensional spacetime and F is an orthonormal n -ad of vectors $\{\xi_1, \dots, \xi_n\}$ at some point $p \in M$. We say an n -dimensional framed spacetime (M', g'_{ab}, F') is a *framed extension* of the n -dimensional framed spacetime (M, g_{ab}, F) if there is a proper isometric embedding $\varphi : M \rightarrow M'$ which takes F into F' .

For each point $p \in M$, the metric assigns a cone structure to the tangent space M_p . Any tangent vector ξ^a in M_p will be *timelike* if $g_{ab}\xi^a\xi^b < 0$, *null* if $g_{ab}\xi^a\xi^b = 0$, or *spacelike* if $g_{ab}\xi^a\xi^b > 0$. Null vectors create the cone structure; timelike vectors are inside the cone while spacelike vectors are outside. A *time orientable* spacetime is one that has a continuous timelike vector field on M . In what follows, it is assumed that spacetimes are time orientable.

For some connected interval $I \subseteq \mathbb{R}$, a smooth curve $\gamma : I \rightarrow M$ is *timelike* if the tangent vector ξ^a at each point in $\gamma[I]$ is timelike. Similarly, a curve is *null* (respectively, *spacelike*) if its tangent vector at each point is null (respectively, spacelike). A curve is *causal* if its tangent vector at each point is either null or timelike. A causal curve is *future directed* if its tangent vector at each point falls in or on the future lobe of the light cone. We say a timelike curve $\gamma : [s, s'] \rightarrow M$ is *closed* if $\gamma(s) = \gamma(s')$. A spacetime (M, g_{ab}) satisfies *chronology* if it does not contain a closed timelike curve. For any two points $p, q \in M$, we write $p \ll q$ if there exists a future-directed timelike curve from p to q . This relation allows us to define the *timelike past* of a point p : $I^-(p) = \{q : q \ll p\}$. We say a spacetime (M, g_{ab}) satisfies *past distinguishability* if there do not exist distinct points $p, q \in M$ such that $I^-(p) = I^-(q)$. We say a set $S \subset M$ is *achronal* if there do not exist $p, q \in S$ such that $p \in I^-(q)$.

An *extension* of a curve $\gamma : I \rightarrow M$ is a curve $\gamma' : I' \rightarrow M$ such that I is a proper subset of I' and $\gamma(s) = \gamma'(s)$ for all $s \in I$. A curve is *maximal* if it has no extension. A curve $\gamma : I \rightarrow M$ in a spacetime (M, g_{ab}) is a *geodesic* if $\xi^a \nabla_a \xi^b = \mathbf{0}$ where ξ^a is the tangent vector and ∇_a is the unique derivative operator compatible with g_{ab} . A point $p \in M$ is a *future endpoint* of a future directed causal curve $\gamma : I \rightarrow M$ if, for every neighborhood O of p , there exists a point $t_0 \in I$ such that $\gamma(t) \in O$ for all $t > t_0$. A *past endpoint* is defined similarly. A causal curve is *inextendible* if it has no future or past endpoint. A causal geodesic $\gamma : I \rightarrow M$ in a spacetime (M, g) is *past incomplete* if it is maximal and there is an $r \in \mathbb{R}$ such that $r < s$ for all $s \in I$.

For any set $S \subseteq M$, we define the *past domain of dependence* of S , written $D^-(S)$, to be the set of points $p \in M$ such that every causal curve with past endpoint p and no future endpoint intersects S . The *future domain of dependence* of S , written $D^+(S)$, is defined analogously. The

entire *domain of dependence* of S , written $D(S)$, is just the set $D^-(S) \cup D^+(S)$. The *edge* of an achronal set $S \subset M$ is the collection of points $p \in S$ such that every open neighborhood O of p contains a point $q \in I^+(p)$, a point $r \in I^-(p)$, and a timelike curve from r to q which does not intersect S . A set $S \subset M$ is a *slice* if it is closed, achronal, and without edge. A spacetime (M, g_{ab}) which contains a slice S such that $D(S) = M$ is said to be *globally hyperbolic*.

Given a spacetime (M, g_{ab}) , let T_{ab} be defined by $\frac{1}{8\pi}(R_{ab} - \frac{1}{2}Rg_{ab})$ where R_{ab} is the Ricci tensor and R the scalar curvature associated with g_{ab} . We say that (M, g_{ab}) satisfies the *weak energy condition* if, for each timelike vector ξ^a , we have $T_{ab}\xi^a\xi^b \geq 0$. We say that (M, g_{ab}) is a *vacuum solution* if $T_{ab} = \mathbf{0}$.

Let S be a set. A relation \leq on S is a *partial order* if, for all $a, b, c \in S$: (i) $a \leq a$, (ii) if $a \leq b$ and $b \leq c$, then $a \leq c$, and (iii) if $a \leq b$ and $b \leq a$, then $a = b$. If \leq is a partial ordering on a set S , we say a subset $T \subseteq S$ is *totally ordered* if, for all $a, b \in T$, either $a \leq b$ or $b \leq a$. Let \leq be a partial ordering on S and let $T \subseteq S$ be totally ordered. An *upper bound* for T is an element $u \in S$ such that for all $a \in T$, $a \leq u$. A *maximal element* of S is an element $m \in S$ such that for all $c \in S$, if $m \leq c$, then $c = m$. Zorn's lemma (which is equivalent to the axiom of choice) is the following: Let \leq be a partial order on S . If each totally ordered subset $T \subseteq S$ has an upper bound, there is a maximal element of S .

3 Definition

Recall the standard definition of spacetime inextendibility.

Definition. A spacetime (M, g_{ab}) is *inextendible* if there does not exist a spacetime (M', g'_{ab}) such that there is a proper isometric embedding $\varphi : M \rightarrow M'$.

The definition requires that an inextendible spacetime be “as large as it can be” in the sense that one compares it to a background class of all “possible” spacetimes. Standardly, one takes this class to be the set of all (smooth, Hausdorff) Lorentzian manifolds as defined in the previous section. But what *should* this class be? The answer is unclear.

Consider Misner spacetime (Hawking and Ellis 1973). Let Misner* be the globally hyperbolic “bottom half” of Misner spacetime. By the standard definition of inextendibility, Misner* is extendible and cannot be extended and remain globally hyperbolic (see below). But suppose that a version of the cosmic censorship conjecture is correct and all physically reasonable spacetimes are globally hyperbolic (Penrose 1979). Then shouldn't Misner* be considered “as large as it can be”? It follows that whether or not Misner* spacetime should count as inextendible depends crucially on the outcome of this version of the cosmic censorship conjecture – a conjecture which is far from settled (Earman 1995, Penrose 1999) and perhaps may never be settled (Manchak, 2011).

Because of examples like these, one is tempted to revise the definition

of inextendibility.² But Geroch (1970, 278) has argued that a revision is less urgent if one can show, for a variety of physically reasonable properties \mathcal{P} , that the following is true.

(*) Every \mathcal{P} -inextendible \mathcal{P} -spacetime is inextendible.

The significance of (*) is this: If a property \mathcal{P} satisfies (*), then any \mathcal{P} -spacetime is inextendible if and only if it is \mathcal{P} -inextendible. Effectively, it makes no difference in such cases whether one defines inextendibility relative to the standard class of all spacetimes or a revised class of all \mathcal{P} -spacetimes. Accordingly, one would like to investigate (*) with respect to a variety of properties \mathcal{P} . Already from the Misner* example above, we have the following (a proof is provided in the Appendix).

Proposition 1. If \mathcal{P} is global hyperbolicity, (*) is false.

Are there physically reasonable properties \mathcal{P} which render (*) true? Geroch (1970, 289) has suggested a number of good candidates including: being a vacuum solution, satisfying chronology, and satisfying an energy condition. The first two cases are still open. Here, we settle the case where \mathcal{P} is the weak energy condition (a proof is provided in the Appendix).

Proposition 2. If \mathcal{P} is the weak energy condition, (*) is false.

We see that the prospect of avoiding the need to revise to the definition of inextendibility does not look good. In the meantime, we may conclude that it is not yet clear that the standard definition captures the intuitive idea that an inextendible spacetime is “as large as it can be”.

4 Metaphysics

A number of experts in general relativity (Penrose 1969, Geroch 1970, Clarke 1976), seem to be committed to the following.

“Metaphysical considerations suggest that to be a serious candidate for describing actuality, a spacetime should be [inextendible]. For example, for the Creative Force to actualize a proper subpart of a larger spacetime would seem to be a violation of Leibniz’s principles of sufficient reason and plenitude. If one adopts the image of spacetime as being generated or built up as time passes then the dynamical version of the principle of sufficient reason would ask why the Creative Force would stop building if it is possible to continue” (Earman 1995, 32).

These metaphysical views are underpinned by an important result due to Geroch (1970).

²See Manchak (2016a) for an extended discussion.

Proposition 3. Every extendible spacetime has an inextendible extension.

The result (which makes use of Zorn’s lemma) seems to show that the Creative Force can always build spacetime until it is no longer possible to build. But of course, this interpretation presupposes that we have been working with the proper definition of inextendibility. And as we have noted, it is not yet clear that we are. Accordingly, one would like to know, for a variety of physically reasonable properties \mathcal{P} , whether the following version of the Geroch (1970) result is true.

(**) Every \mathcal{P} -extendible \mathcal{P} -spacetime has a \mathcal{P} -inextendible \mathcal{P} -extension.

With a bit of work (and Zorn’s lemma), one can show the following (a proof is provided in the Appendix).

Proposition 4. If \mathcal{P} is chronology, (**) is true.

We see that if we revise the definition of inextendibility to be relative to the class of all chronological spacetimes (rather than the standard class of all spacetimes), we have an analogue of the Geroch (1970) result. This is certainly good news for those committed to the metaphysical views expressed above. But are there physically reasonable properties \mathcal{P} which render (**) false? There are.

Of course, spacetime properties may be considered physically reasonable in various senses. Let us conservatively restrict attention a property usually taken to be satisfied by models of our own universe: the property of having every inextendible timelike geodesic be past incomplete. Let us call this the *big bang property* given that it is satisfied by all of the standard “big bang” cosmological models. We are now in a position to state the following (Manchak 2016b).

Proposition 5. If \mathcal{P} is the big bang property, (**) is false.

We see that if we revise the definition of inextendibility to be relative to the class of all spacetimes with the big bang property (rather than the standard class of all spacetimes), we do not have an analogue of the Geroch (1970) result. It is not yet clear that the Creative Force always has the option of building spacetime to be “as large as it can be”.

5 Epistemology

What observational evidence is there (or could there be) in support of the position that spacetime is “as large as it can be”? Following Malament (1977), let us say that a spacetime (M, g_{ab}) is *observationally indistinguishable* from another spacetime (M', g'_{ab}) if, for every point $p \in M$, there is a point $p' \in M'$ such that $I^-(p)$ and $I^-(p')$ are isometric. One can show the following (Manchak 2011).

Proposition 6. Every chronological spacetime is observationally indistinguishable from some other (non-isometric) spacetime which is extendible.

Under the standard definition of inextendibility, it seems that any observer in a chronological spacetime is not in a position to know that her spacetime is “as large as it can be”. But this interpretation presupposes that we have been working with the proper definition of inextendibility. And as we have noted, it is not yet clear that we are. Accordingly, one would like to know, for a variety of physically reasonable properties \mathcal{P} , whether the following version of the Manchak (2011) result is true.

(***) Every chronological \mathcal{P} -spacetime is observationally indistinguishable from some other (non-isometric) \mathcal{P} -spacetime which is \mathcal{P} -extendible.

It turns out that a large class of physically reasonable properties render (***) true. Following Manchak (2011), let us say that a property \mathcal{P} on a spacetime is *local* if, given any two locally isometric spacetimes (M, g_{ab}) and (M', g'_{ab}) , (M, g_{ab}) has \mathcal{P} if and only if (M', g'_{ab}) has \mathcal{P} . Local properties include being a vacuum solution and satisfying the weak energy condition. We are now in a position to state the following (a proof is provided in the Appendix).

Proposition 7. If \mathcal{P} is a local property, (***) is true.

We see that, whenever \mathcal{P} is a local property, if we revise the definition of inextendibility to be relative to the class of all \mathcal{P} -spacetimes (rather than the standard class of all spacetimes), we have an analogue of the Manchak (2011) result. It is not yet clear that we can ever have observational evidence that spacetime is “as large as it can be”.

6 Conclusion

In the preceding, we have registered some skepticism with respect to the position that spacetime must be inextendible – that it must be “as large as it can be” in some sense. We have done this in a variety of ways. First we have shown that it is not yet clear that the standard definition of inextendibility captures the intuitive idea that an inextendible spacetime is “as large as it can be”. Second we have shown, by exploring some plausible revisions to the definition of inextendibility, that it is not yet clear that a spacetime can always be extended to be “as large as it can be”. Finally we have shown, by exploring a class of plausible revisions to the definition of inextendibility, that it is not yet clear that we can ever know that spacetime is “as large as it can be”.

7 Appendix

Proposition 1. If \mathcal{P} is global hyperbolicity, (*) is false.

Proof. Let (N, g_{ab}) be Misner spacetime. So, $N = \mathbb{R} \times \mathbb{S}$ and $g_{ab} = 2\nabla_{(a}t\nabla_{b)}\varphi - t\nabla_a\varphi\nabla_b\varphi$ where the points (t, φ) are identified with the points $(t, \varphi + 2\pi n)$ for all integers n . Now, let $M = \{(t, \varphi) \in N : t < 0\}$ and consider the spacetime (M, g_{ab}) . Clearly, it is extendible. It is also globally hyperbolic since the slice $S = \{(t, \varphi) \in M : t = -1\}$ is such that $D(S) = M$. We need only show that any extension to (M, g_{ab}) fails to be globally hyperbolic.

Let (M', g'_{ab}) be any extension of (M, g_{ab}) and let p be a point in $\partial M \cap M'$. In any neighborhood of p , there will be a point $q \in \partial M \cap M'$ such that $q \neq p$. One can verify that $I^-(p) = M = I^-(q)$. Thus, (M', g'_{ab}) is not past distinguishing and therefore not globally hyperbolic (Hawking and Ellis 1973). \square

Proposition 2. If \mathcal{P} is the weak energy condition, (*) is false.

Proof. Consider Minkowski spacetime $(\mathbb{R}^4, \eta_{ab})$ in standard (t, x, y, z) coordinates where $\eta_{ab} = -\nabla_a t \nabla_b t + \nabla_a x \nabla_b x + \nabla_a y \nabla_b y + \nabla_a z \nabla_b z$. Let $\Omega : \mathbb{R}^4 \rightarrow \mathbb{R}$ be the function defined by $\Omega(t, x, y, z) = \exp(t^3)$. Consider the spacetime (\mathbb{R}^4, g_{ab}) where $g_{ab} = \Omega^2 \eta_{ab}$. Associated with g_{ab} we have (Wald 1984, 446):

$$\begin{aligned} R_{ab} &= -2\nabla_a \nabla_b t^3 - \eta_{ab} \eta^{cd} \nabla_c \nabla_d t^3 + 2(\nabla_a t^3)(\nabla_b t^3) \\ &\quad - 2\eta_{ab} \eta^{cd} (\nabla_c t^3)(\nabla_d t^3) \\ R &= \frac{1}{\Omega^2} [-6\eta^{ab} \nabla_a \nabla_b t^3 - 6\eta^{ab} (\nabla_a t^3)(\nabla_b t^3)]. \end{aligned}$$

We note that $\nabla_a t^3 = 3t^2 \nabla_a t$ and $\nabla_a \nabla_b t^3 = 6t(\nabla_a t)(\nabla_b t)$. Of course, $\eta^{ab}(\nabla_a t)(\nabla_b t) = -1$. Simplifying, we have:

$$\begin{aligned} R_{ab} &= (18t^4 - 12t)(\nabla_a t)(\nabla_b t) + (18t^4 + 6t)\eta_{ab} \\ R &= \frac{1}{\Omega^2} [36t + 54t^4]. \end{aligned}$$

Einstein's equation $R_{ab} - \frac{1}{2}Rg_{ab} = 8\pi T_{ab}$ requires that:

$$T_{ab} = \frac{1}{8\pi} [(18t^4 - 12t)(\nabla_a t)(\nabla_b t) - (9t^4 + 12t)\eta_{ab}]$$

In (\mathbb{R}^4, g_{ab}) , consider any timelike vector $\xi^a = k_0(\partial/\partial t)^a + k_1(\partial/\partial x)^a + k_2(\partial/\partial y)^a + k_3(\partial/\partial z)^a$ where $k_0, k_1, k_2, k_3 \in \mathbb{R}$ and $k_0^2 > k_1^2 + k_2^2 + k_3^2$. We have:

$$\begin{aligned} T_{ab}\xi^a\xi^b &= \frac{1}{8\pi} [(18t^4 - 12t)k_0^2 + (9t^4 + 12t)(k_0^2 - k_1^2 - k_2^2 - k_3^2)] \\ &= \frac{1}{8\pi} [t^4(27k_0^2 - 9(k_1^2 + k_2^2 + k_3^2)) - 12t(k_1^2 + k_2^2 + k_3^2)] \end{aligned}$$

Because $k_0^2 > k_1^2 + k_2^2 + k_3^2$, we know $t^4(27k_0^2 - 9(k_1^2 + k_2^2 + k_3^2)) \geq 0$. And for $t \leq 0$, we know $-12t(k_1^2 + k_2^2 + k_3^2) \geq 0$. It follows that $T_{ab}\xi^a\xi^b \geq 0$ for $t \leq 0$.

Let $M = \{(t, x, y, z) \in \mathbb{R}^4 : t < 0\}$. We have shown that the spacetime $(M, g_{ab|_M})$ is such that it satisfies the weak energy condition and is extendible. It remains for us to show that any extension to $(M, g_{ab|_M})$ fails to satisfy the weak energy condition.

Let (M', g'_{ab}) be any extension to $(M, g_{ab|_M})$. Let p be a point in $\partial M \cap M'$. Let (O, φ) be a chart with $p \in O$ such that we can extend the coordinates (t, x, y, z) on M to $M \cup O \subset M'$. So, for some $p_1, p_2, p_3 \in \mathbb{R}$ we have $p = (0, p_1, p_2, p_3)$. Find some $\delta > 0$ such that $(\delta, p_1, p_2, p_3) \in O$. For $t \in (-\delta, \delta)$, let $p(t) = (t, p_1, p_2, p_3) \in M'$.

Consider the smooth function $f : (-\delta, \delta) \rightarrow \mathbb{R}$ given by $f(t) = g'_{ab}\zeta^a\zeta^b|_{p(t)}$ where $\zeta^a = \sqrt{2}(\partial/\partial t)^a + (\partial/\partial x)^a$. Of course, for all $t < 0$, we have:

$$g'_{ab}\zeta^a\zeta^b = g_{ab}\zeta^a\zeta^b = \Omega^2\eta_{ab}\zeta^a\zeta^b = -\Omega^2 = -\exp(2t^3)$$

Smoothness requires that $f(0) = -1$. This allows us to find an $\epsilon \in (0, \delta)$ such that $f(t) < 0$ for all $t \in (-\epsilon, \epsilon)$. So ζ^a is timelike at $p(t)$ for all $t \in (-\epsilon, \epsilon)$.

Consider the smooth function $g : (-\epsilon, \epsilon) \rightarrow \mathbb{R}$ given by $g(t) = T'_{ab}\zeta^a\zeta^b|_{p(t)}$ where T'_{ab} is defined on M' in the natural way (using the metric g'_{ab} and Einstein's equation). Of course, for all $t < 0$, we have:

$$T'_{ab} = T_{ab} = \frac{1}{8\pi}[(18t^4 - 12t)(\nabla_a t)(\nabla_b t) - (9t^4 + 12t)\eta_{ab}]$$

Because $\nabla_a t \nabla_b t \zeta^a \zeta^b = 2$ and $\eta_{ab}\zeta^a\zeta^b = -1$, we have for all $t < 0$:

$$T'_{ab}\zeta^a\zeta^b = T_{ab}\zeta^a\zeta^b = \frac{1}{8\pi}[(36t^4 - 24t) + (9t^4 + 12t)] = \frac{1}{8\pi}[45t^4 - 12t]$$

Smoothness requires that $g(0) = 0$ and $\frac{d}{dt}g(0) = -\frac{3}{2\pi}$. This allows us to find a $\gamma \in (0, \epsilon)$ such that $g(t) < 0$ for $t \in (0, \gamma)$. Thus, the weak energy condition is violated at $p(t)$ for all $t \in (0, \gamma)$. \square

Definition. Let \mathfrak{F} denote the set of framed spacetimes. Let \leq denote the relation on \mathfrak{F} such that $(M, g_{ab}, F) \leq (M', g'_{ab}, F')$ iff (M', g'_{ab}, F') is a framed extension of (M, g_{ab}, F) .

Lemma 1. The relation \leq is a partial ordering on \mathfrak{F} .

Proof. See Geroch (1969, 188-189). \square

Lemma 2. Let \mathfrak{C} denote the set of framed spacetimes which satisfy chronology. \mathfrak{C} is partially ordered by \leq . Every subset $\mathfrak{T} \subset \mathfrak{C}$ which is

totally ordered by \leq has an upper bound in \mathfrak{C} .

Proof. Since $\mathfrak{C} \subset \mathfrak{F}$, it follows from Lemma 1 that \mathfrak{C} is partially ordered by \leq . Let $\mathfrak{T} = \{(M_i, g_i, F_i)\}$ be a subset of \mathfrak{C} which is totally ordered by \leq . Following Hawking and Ellis (1973, 249), let M be the union of all the M_i where, for $(M_i, g_i, F_i) \leq (M_j, g_j, F_j)$, each $p_i \in M_i$ is identified with $\varphi_{ij}(p_i)$ where $\varphi_{ij} : M_i \rightarrow M_j$ is the unique isometric embedding which takes F_i into F_j . The manifold M will have an induced metric g equal to $\varphi_{i*}g_i$ on each $\varphi_i[M_i]$ where $\varphi_i : M_i \rightarrow M$ is the natural isometric embedding. Finally, take F to be the result of carrying along a chosen F_i using $\varphi_i : M_i \rightarrow M$. Consider the framed spacetime (M, g, F) . We claim it is an upper bound for \mathfrak{T} . Clearly, for all i , we have $(M_i, g_i, F_i) \leq (M, g, F)$. We need only show that $(M, g, F) \in \mathfrak{C}$.

Suppose $(M, g, F) \notin \mathfrak{C}$ and let $\gamma \subset M$ be (the image of) a closed time-like curve. As a topological space (with induced topology from M), γ is compact. For all i , let γ_i be $\gamma \cap M_i$. So, $A = \{\gamma_i\}$ is an open cover of γ . By compactness, there must be a finite subset $A' \subset A$ which is also a cover of γ . One can use the relation \leq on \mathfrak{T} to order the finite number of elements in A' into a nested sequence of subsets $\gamma_j \subseteq \dots \subseteq \gamma_k$. It follows that $\gamma_k = \gamma$. So, $(M_k, g_k, F_k) \notin \mathfrak{C}$: a contradiction. \square

Proposition 4. If \mathcal{P} is chronology, (***) is true.

Proof. Let \mathcal{P} be chronology and let (M, g_{ab}) be a \mathcal{P} -spacetime which is \mathcal{P} -extendible. Let F be an orthonormal n -ad at some point $p \in M$. So, $(M, g_{ab}, F) \in \mathfrak{C}$ where \mathfrak{C} is the set of framed spacetimes which satisfy chronology. By Lemma 2 and Zorn's lemma, there is a maximal element $(M', g'_{ab}, F') \in \mathfrak{C}$ such that $(M, g_{ab}, F) \leq (M', g'_{ab}, F')$. It follows that (M', g'_{ab}) is a \mathcal{P} -inextendible \mathcal{P} -extension of (M, g_{ab}) . \square

Proposition 7. If \mathcal{P} is a local property, (***) is true.

Proof. Let \mathcal{P} be any local property and let (M, g_{ab}) be any chronological \mathcal{P} -spacetime. Now construct (M', g'_{ab}) according to the method outlined in Manchak (2009). Note that (M', g'_{ab}) is a \mathcal{P} -spacetime by construction. Next, remove any point in the $M(1, b)$ portion of the manifold M' and call the resulting spacetime (M'', g''_{ab}) . One can verify that (i) (M, g_{ab}) is observationally indistinguishable from (M'', g''_{ab}) , (ii) (M'', g''_{ab}) is a \mathcal{P} -spacetime, (iii) (M, g_{ab}) is not isometric to (M'', g''_{ab}) . Since (M', g'_{ab}) is a \mathcal{P} -extension to (M'', g''_{ab}) , the latter is \mathcal{P} -extendible. \square

References

- [1] Clarke, C. (1976), "Spacetime Singularities", *Communications in Mathematical Physics*, 49: 17-23.

- [2] Earman, J. (1995). *Bangs, Crunches, Whimpers, and Shrieks: Singularities and Acausalities in Relativistic Spacetimes*. Oxford: Oxford University Press.
- [3] Geroch, R. (1969), "Limits of Spacetimes," *Communications in Mathematical Physics*, 13: 180-193.
- [4] Geroch, R. (1970), "Singularities," in M. Carmeli, S. I. Fickler, and L. Witten (eds.), *Relativity*. New York: Plenum Press, 259-291.
- [5] Geroch, R., and Horowitz, G. (1979). "Global Structure of Spacetimes," in S. W. Hawking and W. Israel (eds), *General Relativity: An Einstein Centenary Survey*. Cambridge: Cambridge University Press, 212-293.
- [6] Hawking, S., and Ellis, G. (1973), *The Large Scale Structure of Space-Time*. Cambridge: Cambridge University Press.
- [7] Malament, D. (1977), "Observationally Indistinguishable Spacetimes," In J. Earman, C. Glymour, & J. Stachel (eds.), *Foundations of Space-Time Theories, Minnesota Studies in the Philosophy of Science Vol. VIII*. Minneapolis: University of Minnesota Press, 61-80.
- [8] Malament, D. (2012). *Topics in the Foundations of General Relativity and Newtonian Gravitation Theory*. Chicago: University of Chicago Press.
- [9] Manchak, J. (2009), "Can We Know the Global Structure of Spacetime?" *Studies in History and Philosophy of Modern Physics*, 40: 53-56.
- [10] Manchak, J. (2011). "What is a Physically Reasonable Spacetime?" *Philosophy of Science*, 78: 410-420.
- [11] Manchak, J. (2016a), "Epistemic 'Holes' in Spacetime." Forthcoming in *Philosophy of Science*.
- [12] Manchak, J. (2016b), "Is the Universe As Large as It Can Be?" Forthcoming in *Erkenntnis*.
- [13] Penrose, R. (1969) "Gravitational Collapse: The Role of General Relativity," *Revisita del Nuovo Cimento*, Serie I, 1: 252-276.
- [14] Penrose, R. (1979). "Singularities and Time-Asymmetry," in S. Hawking and W. Israel (eds), *General Relativity: An Einstein Centenary Survey*. Cambridge: Cambridge University Press, 581-638.
- [15] Penrose, R. (1999), "The Question of Cosmic Censorship", *Journal of Astrophysics and Astronomy*, 20: 233-248.
- [16] Wald, R. (1984), *General Relativity*. Chicago: University of Chicago Press.