

The Motion of a Body in Newtonian Theories¹

James Owen Weatherall²
Logic and Philosophy of Science
University of California, Irvine

Abstract

A theorem due to Bob Geroch and Pong Soo Jang [“Motion of a Body in General Relativity.” *Journal of Mathematical Physics* **16**(1), (1975)] provides the sense in which the geodesic principle has the status of a theorem in General Relativity (GR). Here we show that a similar theorem holds in the context of geometrized Newtonian gravitation (often called Newton-Cartan theory). It follows that in Newtonian gravitation, as in GR, inertial motion can be derived from other central principles of the theory.

1 Introduction

The geodesic principle in General Relativity (GR) states that free massive test point particles traverse timelike geodesics. It has long been believed that, given the other central postulates of GR, the geodesic principle can be proved as a theorem. In our view, though previous attempts³ were highly suggestive, the sense in which the geodesic principle is a theorem of GR was finally clarified by Geroch and Jang (1975).⁴ They proved the following (the statement of which is indebted to Malament (2010, Prop. 2.5.2)):

Theorem 1.1 (Geroch and Jang, 1975) *Let (M, g_{ab}) be a relativistic spacetime, with M orientable. Let $\gamma : I \rightarrow M$ be a smooth, imbedded curve. Suppose that given any open subset O of M containing $\gamma[I]$, there exists a smooth symmetric field T^{ab} with the following properties.*

1. T^{ab} satisfies the strict dominant energy condition, *i.e.* given any future-directed time-like covectors ξ_a, η_a at any point in M , either $T^{ab} = \mathbf{0}$ or $T^{ab}\xi_a\eta_b > 0$;
2. T^{ab} satisfies the conservation condition, *i.e.* $\nabla_a T^{ab} = \mathbf{0}$;

¹I am indebted to David Malament for helpful comments on previous drafts of this paper, and for suggesting the topic. Thank you, too, to helpful audiences in Paris and Wuppertal, and particularly to Harvey Brown and David Wallace.

²weatherj@uci.edu

³For instance, Einstein et al. (1938); Thomas (1962); Taub (1962); Dixon (1964) as well as references in Geroch and Jang (1975).

⁴See also Ehlers and Geroch (2004), who prove a version of the Geroch-Jang theorem that permits backreaction of the particle on the metric.

3. $\text{supp}(T^{ab}) \subset O$; and

4. there is at least one point in O at which $T^{ab} \neq \mathbf{0}$.

Then γ is a timelike curve that can be reparametrized as a geodesic.

The interpretation of the Geroch-Jang theorem can be put as follows: if γ is a smooth curve about which it is possible to construct an arbitrarily small matter field satisfying the conservation and strict dominant energy conditions, then γ can be reparametrized as a timelike geodesic. More roughly, the only curves about which matter can propagate are timelike geodesics. The Geroch-Jang approach has many virtues that previous attempts lacked: (1) Geroch and Jang do not make any specific assumptions about the kinds of matter fields that might compose the free massive test point particle (i.e. they do not need to assume it is a perfect fluid or a dust, etc.), aside from general assumptions that *any* body in GR would be expected to satisfy; (2) Geroch and Jang are able to show that a free massive test point particle traverses a curve *within spacetime*, as opposed to a “line singularity” (cf. Einstein et al., 1938); and (3) Geroch and Jang do not need to make simplifying assumptions regarding the mass multi-pole structure of their test objects (cf. Dixon, 1964).

In so-called “geometrized Newtonian gravitation,” sometimes known as Newton-Cartan theory, the motion of a free massive test point particle is again governed by a geodesic principle. But thus far, little attention has been paid to the question of whether here, too, the geodesic principle has the status of a theorem.⁵ The central result of the present paper (Theorem 4.4 and Corollaries 4.5 and 4.6) is that a direct parallel to the Geroch-Jang theorem does hold in geometrized Newtonian gravitation.⁶ It is worth noting that in the course of proving the geodesic principle as a theorem of geometrized Newtonian gravitation, we prove

⁵The exception is work by Duval and Künzle (1978), who use variational methods to derive the equations of motion of a particle in geometrized Newtonian gravitation. The present paper should be understood to bear the same relationship to Duval and Künzle’s work that Geroch and Jang (1975) bears to attempts to derive the geodesic principle using variational methods in GR. They represent different routes to (almost) the same place. It is our view that for some purposes, the Geroch-Jang method is considerably more perspicuous.

⁶At least, the Geroch-Jang theorem and Theorem 4.4 of this paper are directly parallel mathematically. There is a second class of questions that one might ask, concerning the interpretations of the two theorems in the contexts of their respective spacetime theories. For instance, one might wonder if the conservation condition is as natural an assumption in geometrized Newtonian gravitation theory as in GR. We do not address this such questions here, but will return to them in future work.

a lemma that can be understood as a proof of Newton's first law (appropriately reformulated in covariant, four dimensional language) in non-geometrized Newtonian gravitation. Thus we show that the principles governing inertial motion in both standard Newtonian theory and geometrized Newtonian gravitation are dependent on the other principles of the theory, just as in GR.

The remainder of the paper will proceed as follows. In section 2 we will briefly review geometrized Newtonian gravitation. Section 3 will establish several important preliminaries concerning integration in classical spacetime, as well as appropriate definitions of momentum flux, angular momentum flux, and center of mass in the geometrized context. The main results of the paper will be presented in section 4.

2 Review of Geometrized Newtonian Gravitation

Geometrized Newtonian gravitation was first developed in the 1920s by Élie Cartan (1923, 1924) and, apparently independently, by Kurt Friedrichs (1927), with substantial later contributions by Ehlers (1981), Künzle (1976), Trautman (1965), and others (see Malament (2010, Ch. 4) for an extensive list of references). Geometrized Newtonian gravitation is distinctive because it re-casts standard Newtonian dynamics and gravitation in a geometrical language, bringing it as close to GR as possible. Indeed, it can be shown that in a precise sense, geometrized Newtonian gravitation is a classical limit of GR (Künzle, 1976; Ehlers, 1981; Malament, 1986). It is thus the ideal context for work that seeks to compare GR with classical Newtonian physics. In this section we will briefly review the central concepts of geometrized Newtonian gravitation. We will not describe the full details of the theory; rather, the focus will be on setting up the language in which we will operate in the remainder of the paper. For details, we recommend Malament (2010, Ch. 4), which is (to our knowledge) the most systematic treatment of the subject available.

We begin by defining a classical spacetime.

Definition 2.1 *A classical spacetime is an ordered quadruple $(M, t_{ab}, h^{ab}, \nabla)$, where M is*

a smooth,⁷ connected, four dimensional manifold; t_{ab} is a smooth symmetric field on M of signature $(1, 0, 0, 0)$; h^{ab} is a smooth symmetric field on M of signature $(0, 1, 1, 1)$; and ∇ is a derivative operator on M compatible with t_{ab} and h^{ab} , i.e. it satisfies $\nabla_a t_{bc} = \nabla_a h^{bc} = \mathbf{0}$. We additionally require that t_{ab} and h^{ab} are orthogonal, i.e. $t_{ab}h^{bc} = \mathbf{0}$.

Note that “signature,” here, has been extended to cover the degenerate case. We can see immediately from the signatures of t_{ab} and h^{ab} that neither is invertible. Hence in general neither t_{ab} nor h^{ab} can be used to raise and lower indices.

The field t_{ab} can be thought of as a temporal metric on M in the sense that given any vector ξ^a in the tangent space at a point, p , $\|\xi^a\| = (t_{ab}\xi^a\xi^b)^{1/2}$ is the *temporal length* of ξ^a at that point. If the temporal length of ξ^a is positive, ξ^a is *timelike*; otherwise, it is *spacelike*. At any point, it is possible to find a covector t_a , unique up to a sign, such that $t_{ab} = t_a t_b$. If there is a continuous, globally defined vector field t_a such that at every point $t_{ab} = t_a t_b$, then the spacetime is *temporally orientable* (we will encode the assumption that a spacetime is temporally oriented by replacing t_{ab} with t_a in our definitions of classical spacetimes). h^{ab} , meanwhile, can be thought of as a spatial metric. However, since there is no way to lower the indices of h^{ab} , we cannot calculate the spatial length of a vector directly. Instead, we rely on the fact that if ξ^a is a spacelike vector (as defined above), then there exists a (non-unique) covector σ_a such that $\xi^a = h^{ab}\sigma_b$. The *spatial length* of ξ^a can then be defined as $(h^{ab}\sigma_a\sigma_b)^{1/2}$. It can be shown that this length is independent of the choice of σ_a . If ξ^a is not a spacelike vector, then there is no way to assign it a spatial length. Note, too, that it is possible to define the Riemann curvature tensor $R^a{}_{bcd}$ and the Ricci tensor R_{ab} with respect to ∇ as in GR (or rather, as in differential geometry generally). Flatness ($R^a{}_{bcd} = \mathbf{0}$) carries over intact from GR; we say a classical spacetime is *spatially flat* if $R^{abcd} = R^a{}_{nmq}h^{bn}h^{cm}h^{dq} = \mathbf{0}$. This latter condition is equivalent to $R^{ab} = h^{an}h^{bm}R_{nm} = \mathbf{0}$.⁸

We describe matter in close analogy with GR. Massive point particles are represented by their worldlines, which are smooth future-directed timelike curves parameterized by elapsed

⁷We will explicitly indicate that various fields are smooth in the statements of lemmas and theorems, but throughout the supporting discussion, we will at times take for granted that any object that is a candidate for smoothness is indeed smooth.

⁸See Malament (2010, Prop. 4.15).

time. (Point particles in the current framework have the same attenuated status as in GR—really, we are thinking of a field theory, and point particles are some appropriate idealization.) For a point particle with mass m , we can always define a smooth unit vector field ξ^a tangent to its worldline (the *four-velocity*), such that we can define a *four-momentum* field, $p^a = m\xi^a$. Thus the mass of the particle is given by the temporal length of its four-momentum. In similar analogy to the relativistic case, we can associate with any matter field a smooth symmetric field T^{ab} . T^{ab} encodes the four-momentum density of the matter field as determined by a future directed timelike observer at a point, but in this case all observers agree on the four-momentum density at any point q : $(p^a)_{|q} = (t_b T^{ab})_{|q}$. Contracting once more with t_b yields the mass density, $\rho = t_a t_b T^{ab}$. Since T^{ab} encodes mass and momentum density in geometrized Newtonian gravitation, rather than energy and momentum density (as in GR), it is called the *mass-momentum tensor*. It is standard to assume that mass density is positive whenever $T^{ab} \neq \mathbf{0}$, i.e. $\rho = T^{ab} t_a t_b > 0$. This condition, called the *mass condition*, takes the place of the various energy conditions in GR.

In the present covariant four dimensional language, standard Newtonian mechanics can be expressed as follows. Let (M, t_a, h^{ab}, ∇) be a classical spacetime. We require that ∇ is flat. We begin by considering the dynamics of a test point particle with mass m and four-velocity ξ^a . The acceleration of the particle’s worldline, $\xi^b \nabla_b \xi^a$, is determined by the external forces acting on the particle according to the relation $F^a = m \xi^b \nabla_b \xi^a$. In the absence of external forces, a massive test point particle undergoes geodesic motion. If the total mass-momentum content of spacetime is described by T^{ab} , we require that the *conservation condition* holds, i.e. at every point $\nabla_a T^{ab} = \mathbf{0}$. To add gravitation to the theory, we can represent the gravitational potential as a smooth scalar field φ on M . φ is required to satisfy Poisson’s equation, $\nabla_a \nabla^a \varphi = 4\pi\rho$ (where ∇^a is shorthand for $h^{ab} \nabla_b$). Gravitation is considered a force; the gravitational force on a point particle is given by $F^a = -m \nabla^a \varphi$.

In geometrized Newtonian gravitation we again begin with a classical spacetime (M, t_a, h^{ab}, ∇) , but now we allow ∇ to be curved. Once again, the acceleration of a particle with mass m

and four-velocity ξ^a is determined by the relation $F^a = m\xi^b\nabla_b\xi^a$, where F^a represents the external forces acting on the particle; likewise, free massive test point particles undergo geodesic motion. However, the geodesics are now determined relative to the not-necessarily-flat derivative operator. The conservation condition is again expected to hold. Gravitation enters the theory via a geometrized form of Poisson’s equation: if T^{ab} describes the total mass-momentum density in the spacetime, then the Ricci curvature tensor $R_{ab} = R^n_{\ abn}$ is given by $R_{ab} = 4\pi\rho t_a t_b$. Since the Riemann curvature tensor (and by extension, the Ricci tensor) is determined by ∇ , the geometrized Poisson’s equation places a constraint on the derivative operator. In particular, ∇ must be such that, for all smooth vector fields ξ^a , $R_{ab}\xi^a = -2\nabla_{[b}\nabla_{n]}\xi^n = 4\pi\rho t_a t_b\xi^a$. Note, too, that the geometrized Poisson’s equation forces spacetime to be spatially flat, because if Poisson’s equation holds, then $R^{ab} = h^{an}h^{bm}R_{nm} = 4\pi\rho h^{an}h^{bm}t_n t_m = \mathbf{0}$ by the orthogonality condition on the metrics.

It is always possible to “geometrize” a gravitational field on a flat classical spacetime—that is, we can always move from the covariant formulation of standard Newtonian gravitation to geometrized Newtonian gravitation, via a result due to Andrzej Trautman (1965).

Proposition 2.2 (Trautman Geometrization Lemma.) (Slightly modified from Malament, 2010, Prop. 4.2.1.) *Let $(M, t_a, h^{ab}, \overset{f}{\nabla})$ be a flat classical spacetime. Let φ and ρ be smooth scalar fields on M satisfying Poisson’s equation, $\overset{f}{\nabla}_a\overset{f}{\nabla}^a\varphi = 4\pi\rho$. Finally, let $\overset{g}{\nabla} = (\overset{f}{\nabla}, C^a_{\ bc})$,⁹ with $C^a_{\ bc} = -t_b t_c \overset{f}{\nabla}^a\varphi$. Then $(M, t_a, h^{ab}, \overset{g}{\nabla})$ is a classical spacetime; $\overset{g}{\nabla}$ is the unique derivative operator on M such that given any timelike curve with (normalized) tangent vector field ξ^a ,*

$$\xi^n \overset{g}{\nabla}_n \xi^a = \mathbf{0} \Leftrightarrow \xi^n \overset{f}{\nabla}_n \xi^a = -\overset{f}{\nabla}^a \varphi; \tag{G}$$

⁹This notation is explained in Malament (2010, Prop. 1.7.3). Briefly, if ∇ is a derivative operator on M , then any other derivative operator on M is determined relative to ∇ by a smooth symmetric (in the lower indices) tensor field, $C^a_{\ bc}$, and so specifying the $C^a_{\ bc}$ field and ∇ is sufficient to uniquely determine a new derivative operator.

and the Riemann curvature tensor relative to $\overset{g}{\nabla}$, $\overset{g}{R}{}^a{}_{bcd}$, satisfies

$$\overset{g}{R}{}_{ab} = 4\pi\rho t_a t_b \tag{CC1}$$

$$\overset{g}{R}{}^a{}_{b\ c} = \overset{g}{R}{}^c{}_{d\ a} \tag{CC2}$$

$$\overset{g}{R}{}^{ab}{}_{cd} = \mathbf{0}. \tag{CC3}$$

Trautmann showed that it is also possible to go in the other direction. That is, given a curved classical spacetime, it is possible to recover a flat classical spacetime and a gravitational field, φ —so long as the curvature conditions (CC1)-(CC3) are met.

Proposition 2.3 (Trautman Recovery Theorem.) (Slightly modified from Malament, 2010, Prop. 4.2.5.) *Let $(M, t_a, h^{ab}, \overset{g}{\nabla})$ be a classical spacetime that satisfies (CC1)-(CC3) for some smooth scalar field ρ . Then, at least locally on M , there exists a smooth scalar field φ and a flat derivative operator on M , $\overset{f}{\nabla}$, such that $(M, t_a, h^{ab}, \overset{f}{\nabla})$ is a classical spacetime; (G) holds for all timelike curves with (normalized) tangent vector field ξ^a ; and φ and $\overset{f}{\nabla}$ together satisfy Poisson's equation, $\overset{f}{\nabla}{}^a \overset{f}{\nabla}{}^a \varphi = 4\pi\rho$.*

It is worth pointing out that the pair $(\overset{f}{\nabla}, \varphi)$ is not unique. It is also worth pointing out that whenever we begin with standard Newtonian theory and move to geometrized Newtonian theory, it is always possible to move back to the standard theory, because Prop. 2.2 guarantees that the curvature conditions (CC1)-(CC3) are satisfied.

3 Some preliminary definitions

We can now proceed to lay the groundwork for the present contribution. Throughout this section, let (M, t_a, h^{ab}, ∇) be a classical spacetime. Let T^{ab} be a smooth symmetric tensor field on M satisfying three conditions: (1) the mass condition, (2) the conservation condition, and (3) given any spacelike hypersurface $\Sigma \subset M$, $\text{supp}(T^{ab}) \cap \Sigma$ is bounded.

For any manifold A , we will denote the space of all smooth tensor fields on A by $\mathfrak{T}(A)$; the space of smooth contravariant fields on A will be $\mathfrak{T}^\bullet(A)$ and the smooth covariant fields on A will be $\mathfrak{T}_\bullet(A)$. Suppose then that $\Sigma \subset M$ is an imbedded submanifold of M . (Note

that we will *always* assume that submanifolds are connected.) The map $\overset{\Sigma}{i} : \Sigma \rightarrow M$ will be assumed to represent the imbedding map (i.e. the identity map); the corresponding pull-back map $\overset{\Sigma}{i}^* : \mathfrak{T}_{\bullet}(M) \rightarrow \mathfrak{T}_{\bullet}(\Sigma)$ represents the restriction of a covariant tensor field on M to a covariant tensor field on Σ . Throughout this section and the next, we will write that a given spacelike hypersurface *slices* the support (or the convex hull, etc.) of T^{ab} . This assertion can be spelled out in a number of ways; one that is adequate for current purposes is as follows. Let $\Sigma \subset M$ be a spacelike hypersurface of M . We will say that Σ slices the support (say) of T^{ab} if and only if $\text{supp}(T^{ab}) \cap \Sigma \neq \emptyset$ and for any spacelike hypersurface $\tilde{\Sigma}$ such that $\Sigma \subseteq \tilde{\Sigma}$, $\text{supp}(T^{ab}) \cap \Sigma = \text{supp}(T^{ab}) \cap \tilde{\Sigma}$. The idea is that there is at least one point $q \in \text{supp}(T^{ab})$ that is also in Σ , and moreover, any points in $\text{supp}(T^{ab})$ that are spacelike related to q are also in Σ .

3.1 Volume Elements in Classical Spacetimes

In what follows, we will make essential use of volume elements on differentiable manifolds with classical spacetime structure. Some work is required to say what is meant by a volume element without a (invertible, non-degenerate) metric in the background. First, the standard notion of orientability carries over intact from more familiar contexts: the underlying manifold of a classical spacetime is *orientable* if it admits a smooth, globally defined, non-vanishing 4-form. In this context, we can define a *volume element* on an orientable manifold as a smooth 4-form ϵ_{abcd} satisfying the normalization condition,

$$\epsilon_{abcd}\epsilon_{efgh}h^{bf}h^{cg}h^{dh} = 6t_a t_e,$$

which is equivalent to requiring that, given any four vectors at any point $p \in M$, if one of them is a unit timelike vector, ξ^a , and the other three are mutually orthogonal unit spacelike vectors, η^a , then $\epsilon_{abcd}\xi^a\eta^1\eta^2\eta^3 = \pm 1$. Dimensionality considerations are sufficient to show that the volume element is unique up to sign. Specifying a volume element on M provides

an orientation for the manifold; when we call a manifold *oriented*, we are assuming a fixed choice of a volume element in the background. Finally, to say two n -forms $\omega_{a_1 \dots a_n}$ and $\omega'_{a_1 \dots a_n}$ are *co-oriented* is to say that $\omega_{a_1 \dots a_n} = f \omega'_{a_1 \dots a_n}$, where $f > 0$ everywhere.

Now we consider hypersurfaces of M . A hypersurface in a classical spacetime is *spacelike* at a point if all of its tangent vectors are; otherwise it is *timelike* at that point. In what follows, we will limit attention to hypersurfaces that are either everywhere spacelike or everywhere timelike. Suppose Σ is a (timelike or spacelike) hypersurface of M . As above, we will say Σ is orientable if it admits a smooth, globally defined, non-vanishing 3-form. Then, if Σ is orientable, it is always possible to factor the volume element on M in the neighborhood of Σ into $\epsilon^M_{abcd} = \tilde{n}^{\Sigma}_{[a} \tilde{\omega}_{bcd]}$, where $\tilde{\omega}_{abc}$ is a (non-unique) 3-form on M and where \tilde{n}_a is a unit covector field normal to Σ . If Σ is spacelike, then $\tilde{n}_a = \pm t_a$; if Σ is timelike, then $h^{ab} \tilde{n}_a \tilde{n}_b = 1$ and whenever $v^a \in \mathfrak{T}^\bullet(M)$ is tangent to Σ , $v^a \tilde{n}_a = 0$. We can then take $\tilde{i}^*(\tilde{\omega}_{abc}) = \tilde{\epsilon}_{abc}$ to define a volume element on Σ (in other words, the restriction to Σ of any 3-form satisfying the factorization condition above gives a volume element on Σ). As above, dimensionality considerations show that volume elements on hypersurfaces are unique up to sign; to say a hypersurface is oriented will be to assume that there's a fixed choice of volume element in the background.

Note that there are in general two possible unit covector fields normal to any given oriented hypersurface of M : if \tilde{n}_a is a unit normal covector field, then so is $-\tilde{n}_a$. However, the sign of \tilde{n}_a as we have defined it is wholly fixed by the relative orientations of M and Σ because ϵ^M_{abcd} is fixed by the orientation of M and the sign of $\tilde{\omega}_{abc}$ is fixed by the orientation of Σ . Thus given any oriented hypersurface of M , there is a unique unit normal covector field that satisfies the stated factorization condition. Conversely, a choice of normal covector field uniquely picks out an orientation for a hypersurface. As a matter of definition, in the special case where Σ is an oriented *spacelike* hypersurface, we will call Σ *future-directed* (relative to the orientation of M) if $\tilde{n}_a = t_a$; likewise, Σ is *past-directed* if $\tilde{n}_a = -t_a$. Finally, if A is an oriented p dimensional manifold, we will denote its volume element by $\epsilon^A_{a_1 \dots a_p}$.

3.2 Integration in Flat Classical Spacetimes

Here and in the next three subsections (§§3.2-3.5), we will assume that ∇ is a flat derivative operator and that M is oriented and simply connected.¹⁰ Under this assumption, we will need to make sense of some improper-looking integrals, in which the integrand and the integral have (the same) contravariant indices. That is, we will consider integrals of the form $\alpha^{a_1 \dots a_n} = \int_S \beta^{a_1 \dots a_n} \omega_{b_1 \dots b_p}$ where S is a three or four dimensional imbedded submanifold of M and ω is a 3- or 4-form, respectively. We make no claims about what such integrals mean (if anything) under general circumstances. However, when ∇ is flat and M is orientable and simply connected, they can be understood as follows. Pick a point, $q \in M$, and let $\{\overset{1}{\sigma}_a(q), \dots, \overset{4}{\sigma}_a(q)\}$ be an orthonormal*¹¹ basis for the cotangent space of M at q .¹² Since ∇ is flat, parallel transport of covectors is (locally) path-independent; since M is simply connected, we can extend the cobasis at q to all points in M without introducing any ambiguities, by parallel transporting each of the cobasis elements to each other point. This method is guaranteed to produce smooth fields of orthonormal covectors on M —that is, fields of constant basis covectors, $\{\overset{1}{\sigma}_a, \dots, \overset{4}{\sigma}_a\}$.¹³

We can define the integrals required in terms of such bases. Taking an integral with a single contravariant index (it is easy to see how to generalize to more indices), we say $\alpha^a = \int_S \beta^a \omega_{b_1 \dots b_p}$ is the vector field such that, given any covector field $\kappa_a \in \mathfrak{T}_\bullet(S)$, $\alpha^a \kappa_a = \sum_{i=1}^4 \overset{i}{\kappa} \overset{i}{\sigma}_a \alpha^a = \sum_{i=1}^4 \overset{i}{\kappa} \int_S \overset{i}{\sigma}_a \beta^a \omega_{b_1 \dots b_p}$, where $\overset{i}{\kappa}$ is defined so that $\kappa_a = \sum_{i=1}^4 \overset{i}{\kappa} \overset{i}{\sigma}_a$.¹⁴ Note that

¹⁰Since any manifold is *locally* simply connected, we can always extend the notion of integral (and likewise, momentum flux, angular momentum flux, and center of mass) described here by limiting attention to simply connected open regions of an arbitrary manifold (construed as submanifolds). We will return to this idea in Corollary 4.5.

¹¹The star indicates that the language is being abused. See Malament (2010, pgs. 168-9).

¹²Nothing rides on the dimensionality of M here, but since we already have a background manifold in place, we are using it for specificity. The construction we are using would work in any space that is flat, orientable, and simply connected.

¹³Why? The field is smooth because parallel transport is smooth. The resulting fields form a basis everywhere because parallel transport preserves temporal and spatial length and the dot product of constant vectors: $\nabla_a (h^{bc} \overset{i}{\sigma}_b \overset{j}{\sigma}_c) = 0$ since all of the relevant fields are, by definition or construction, constant.

¹⁴This definition of the integral is intended to conform to a kind of piecewise integration over the various components of a vector: a generalization of, in the notation of a first-year vector calculus class, $\int_S (\beta_x(x, y, z), \beta_y(x, y, z), \beta_z(x, y, z)) dx dy dz$, which would yield a constant vector.

since S is an imbedded submanifold of M , $\iota^*(\beta^a \sigma_a) = \beta^a \sigma_a \circ \iota = \beta^a \sigma_a$ because $\beta^a \sigma_a$ is a scalar field. The vector α must exist, as the defining relation for the integral generates a map from the covectors to C^∞ . Moreover, it can easily be shown that this definition of the integral is independent of the choice of basis, due to the linearity of the integral.

Finally, it will prove helpful to register up front how to express two well-known facts about integration in the present language.¹⁵ First, suppose that $\Sigma \subset M$ is an oriented, imbedded hypersurface of M and let β^a be an arbitrary contravariant vector field on M . Then we can immediately write $4\beta^a \epsilon_{abcd} = 4\beta^a n_{[a} \omega_{bcd]} = \beta^a n_a \omega_{bcd} - 3n_{[b} \beta^a \omega_{|a|cd]}$. To integrate, we need to take the pull-back to Σ of both sides of this expression, yielding $\iota^*(4\beta^a \epsilon_{abcd}) = \iota^*(\beta^a n_a \omega_{bcd} - 3n_{[b} \beta^a \omega_{|a|cd]}) = \iota^*(\beta^a n_a \omega_{bcd}) = \iota^*(\beta^a n_a) \epsilon_{bcd}$, because the pull-back map commutes with exterior multiplication, and $\iota^*(n_a) = \mathbf{0}$ because n_a is normal to Σ . Thus,

$$\int_{\Sigma} \iota^*(\beta^a \epsilon_{abcd}) = \frac{1}{4} \int_{\Sigma} \iota^*(\beta^a n_a) \epsilon_{bcd}.$$

Secondly, suppose that N is a four dimensional submanifold of M with boundary ∂N , where we assume ∂N can be written as the union of a collection of hypersurfaces, each of which is everywhere timelike or everywhere spacelike. Then if ω_{bcd} is any 3-form on N , we can write Stokes' theorem in the current language as

$$\int_N d_a \omega_{bcd} = \int_N \nabla_{[a} \omega_{bcd]} = \int_{\partial N} \iota^*(\omega_{bcd}),$$

where d represents the exterior derivative on N . Both of these facts will be of use in the ensuing discussion.

¹⁵These are discussed in full rigor by Boothby (2003); we find the lecture notes by van Suijlekom and Hawkins (2009) to be particularly clear, though they are brief.




Figure 1: Σ_1 and Σ_2 slice the support of T^{ab} , which is bounded in any spacelike hypersurface by construction. Σ_3 , then, is a hypersurface that joins Σ_1 and Σ_2 but does not intersect the support of T^{ab} so that $\Sigma_1 \cup \Sigma_2^- \cup \Sigma_3$ forms the boundary of an oriented four dimensional submanifold of M ; here, the arrows give the orientation of S , which is inherited from Σ_1 and Σ_2^- .

3.3 Momentum

We can use this notion of integration in classical spacetimes to define the momentum flux through a spacelike hypersurface.

Definition 3.1 *Given any oriented hypersurface $\Sigma \subset M$, we define the momentum flux through Σ to be $P^a(\Sigma) = \int_{\Sigma} T^{ab} t_b^{\Sigma} \epsilon_{cde} = \int_{\Sigma} p^a \epsilon_{cde}^{\Sigma}$.*

Proposition 3.2 *Let Σ_1, Σ_2 be any two future-directed spacelike hypersurfaces slicing the support of T^{ab} . Then $P^a(\Sigma_1) = P^a(\Sigma_2)$.*

Proof. Let Σ_1 and Σ_2 be two future-directed spacelike hypersurfaces slicing the support of T^{ab} . Consider a third (timelike) hypersurface, Σ_3 , connecting Σ_1 and Σ_2 in such a way that

(1) $\text{supp}(T^{ab}) \cap \Sigma_3 = \emptyset$ and (2) if we reverse the orientation of the temporally prior of the spacelike hypersurfaces (say, Σ_2), then $\partial S \equiv \Sigma_1 \cup \Sigma_2^- \cup \Sigma_3$ forms the boundary of an oriented, simply connected four dimensional submanifold S of M , whose orientation is as given by the normal covectors depicted in Fig. 1. Since the support of T^{ab} does not intersect Σ_3 , it follows immediately that $\int_{\Sigma_3} T^{ab} n_b \epsilon_{cde}^{\Sigma_3} = \mathbf{0}$. Let κ_a be an arbitrary covector field on M . Then,

$$\begin{aligned}
\kappa_a(P^a(\Sigma_1) - P^a(\Sigma_2)) &= \sum_{i=1}^4 \kappa^i \left(\int_{\Sigma_1} T^{ab} \sigma_a t_b \epsilon_{cde}^{\Sigma_1} - \int_{\Sigma_2} T^{ab} \sigma_a t_b \epsilon_{cde}^{\Sigma_2} \right) \\
&= \sum_{i=1}^4 \kappa^i \left(\int_{\Sigma_1} T^{ab} \sigma_a t_b \epsilon_{cde}^{\Sigma_1} + \int_{\Sigma_2^-} T^{ab} \sigma_a t_b \epsilon_{cde}^{\Sigma_2^-} + \int_{\Sigma_3} T^{ab} \sigma_a n_b \epsilon_{cde}^{\Sigma_3} \right) \\
&= \sum_{i=1}^4 \kappa^i \left(\int_{\Sigma_1} \iota^{\Sigma_1} *(T^{ab} \sigma_a t_b) \epsilon_{cde}^{\Sigma_1} + \int_{\Sigma_2^-} \iota^{\Sigma_2^-} *(T^{ab} \sigma_a t_b) \epsilon_{cde}^{\Sigma_2^-} \right. \\
&\quad \left. + \int_{\Sigma_3} \iota^{\Sigma_3} *(T^{ab} \sigma_a n_b) \epsilon_{cde}^{\Sigma_3} \right) = \sum_{i=1}^4 \kappa^i \left(\int_{\partial S} \partial^S \iota^i *(T^{ab} \sigma_a n_b) \epsilon_{cde}^S \right) \\
&= 4 \sum_{i=1}^4 \kappa^i \left(\int_{\partial S} \partial^S \iota^i *(T^{ab} \sigma_a \epsilon_{bcde}^S) \right) = 4 \sum_{i=1}^4 \kappa^i \left(\int_S \nabla_{[n} T^{ab} \sigma_{|a} \epsilon_{b|cde]}^S \right)
\end{aligned}$$

The third equality follows because $T^{ab} t_a \sigma_b$ is a scalar field, and so it is unaffected by the pull-backs; the fifth equality makes use of the relation cited above concerning flux integrals; and the final equality follows by Stokes' theorem.

Consider the integrand of the last of the expressions above, $\nabla_{[n} T^{ab} \sigma_{|a} \epsilon_{b|cde]}^S$, which is a 4-form. The space of n -forms on any n dimensional manifold is one dimensional, and so it must be that $\nabla_{[n} T^{ab} \sigma_{|a} \epsilon_{b|cde]}^S = f \epsilon_{ncde}^S$, for some scalar field f . The goal is to show that f must be zero; if this is the case, then the integrand vanishes. Let ϵ^{abcd} (with raised indices) be a totally anti-symmetric contravariant tensor, normalized so that $\epsilon_{abcd}^S \epsilon^{efgh} = 4! \delta_a^{[e} \delta_b^f \delta_c^g \delta_d^{h]}$. This field can be constructed out of any (contravariant) basis fields for S . Multiplying the

integrand by ϵ^{abcd} and contracting, then, we find

$$\begin{aligned} f \epsilon_{ncde}^S \epsilon^{ncde} &= 4! f = \nabla_{[n} (T^{ab} \sigma_{|a}^i \epsilon_{b|cde]}^S) \epsilon^{ncde} \\ &= 4! \nabla_n (T^{ab} \sigma_a^i) \delta_b^n = 4! \nabla_b T^{ab} \sigma_a^i = 0, \end{aligned}$$

where the last step follows from the conservation condition on T^{ab} . Thus $f = 0$. It follows immediately that $\kappa_a (P^a(\Sigma_1) - P^a(\Sigma_2)) = 0$. But κ_a was an arbitrary covector, which means that $P^a(\Sigma_1) - P^a(\Sigma_2)$ must vanish identically, and so $P^a(\Sigma_1) = P^a(\Sigma_2)$. \square

We have not given an interpretation to T^{ab} yet; however, it is worth noting that if T^{ab} is understood as the Newtonian mass-momentum tensor, Prop. 3.2 is a statement of conservation of momentum. To see why, note that if Σ_1 and Σ_2 are spacelike hypersurfaces slicing the support of T^{ab} , then the momentum flux is the same through both of them. Since we have assumed that M is simply connected, it is possible to define a global time function on the spacetime, and so Prop. 3.2 implies that P^a is constant in time.¹⁶

Definition 3.3 *Let $\Sigma \subset M$ be any spacelike hypersurface slicing the support of T^{ab} . Then the total momentum of the system can be defined pointwise as follows. At any point $p \in M$, $(P^a)_{|p} = P^a(\Sigma)$. By Prop. 3.2, P^a is independent of the choice of surface.*

Proposition 3.4 *The covariant derivative of P^a is given by $\nabla_n P^a = 0$.*

Proof. Fix $o \in M$ and let $\Sigma \subset M$ be any spacelike hypersurface slicing the support of T^{ab} . Then $(P^a)_{|o} = \int_{\Sigma} T^{ab} t_b \epsilon_{cde}^{\Sigma}$. Let $(P^a)_{\parallel o}$ represent the vector field found by parallel transporting $(P^a)_{|o}$ to all points of M . Now take an arbitrary point $p \in M$. By definition, we have $(P^a)_{|p} = \int_{\Sigma} T^{ab} t_b \epsilon_{cde}^{\Sigma} = ((P^a)_{\parallel o})_{|p}$. But p was arbitrary and so $P^a = (P^a)_{\parallel o}$. Since $(P^a)_{\parallel o}$ is constant by construction, P^a must be constant as well. We can conclude that $\nabla_n P^a = 0$. \square

¹⁶In general, it is always possible to define local time functions on a classical spacetime. If we allow M to be non-simply connected, we can limit attention to simply connected open regions of M . We can then calculate momentum flux within the simply connected region, in which case, so long as the local simultaneity slices associated with a given local time function slice the support of T^{ab} , the local P^a will be constant relative to the local time function.

Remark 3.5 Note that P^a is timelike, as $P^a t_a = \int_{\Sigma} T^{ab} t_a t_b \epsilon_{cde}^{\Sigma} > 0$.

3.4 Angular Momentum

We have shown that P^a is a constant timelike vector field relative to ∇ . Thus its integral curves are geodesics. We will work with a normalized vector field, V^a , given by $V^a = P^a / (P^n t_n)$, whose integral curves are also geodesics. In what follows, let Γ be the set of maximal integral curves of V^a . Since ∇ is flat, we can define a class of vector fields, $\{\overset{p}{\chi}^a | p \in M\} \subset \mathfrak{T}^\bullet$, satisfying the following properties: for any $p \in M$, $(\overset{p}{\chi}^a)|_p = \mathbf{0}$ and $\nabla_a \overset{p}{\chi}^b = \delta_a^b$.¹⁷ These can be thought of as fields of “position vectors” centered at a specified point. At each point q , $(\overset{p}{\chi}^a)|_q$ gives the vector “from p to q ” in the tangent space at q .

Definition 3.6 Given any point $p \in M$ and any oriented hypersurface $\Sigma \subset M$, we define the angular momentum flux through Σ relative to p to be $J^{ab}(\Sigma, p) = \int_{\Sigma} \overset{p}{\chi}^{[a} T^{b]c} t_c \epsilon_{def}^{\Sigma}$.

Proposition 3.7 Let Σ_1, Σ_2 be any two future-directed spacelike hypersurfaces slicing the support of T_{ab} and let $p \in M$. Then $J^{ab}(\Sigma_1, p) = J^{ab}(\Sigma_2, p)$.

We omit the proof of this claim, as it follows by identical reasoning as the proof of Prop. 3.2.

Prop. 3.7 is analogous to Prop. 3.2 and can similarly be interpreted as a statement of the conservation of angular momentum about any given point. It justifies a definition analogous to that of P^a .

Definition 3.8 Let $\Sigma \subset M$ be any spacelike hypersurface slicing the support of T^{ab} . Then the total angular momentum, J^{ab} , can be defined pointwise in the following way. At any point $p \in M$, $(J^{ab})|_p = J^{ab}(\Sigma, p)$. By Prop. 3.7, J^{ab} at any point is independent of the choice of Σ .

¹⁷Such a vector field, relative to ∇ , exists everywhere whenever ∇ is flat and the underlying space is orientable and simply connected. To see why, note that at any point p , one can always pick a basis for the tangent vector space at p by taking the tangent vectors at p of a set of coordinate curves through p (see the discussion in Malament (2010, pg. 53)). Since ∇ is flat, one can parallel transport this basis to find a set of coordinate basis fields everywhere. Then the reasoning in the proof of Prop. 1.7.11 of Malament (2010) applies, using the coordinate maps and basis vectors that we have just described.

Proposition 3.9 *The covariant derivative of J^{ab} is given by $\nabla_a J^{bc} = -\delta_a^{[b} P^{c]}$.*

Proof. Fix $o \in M$ and consider any $p \in M$ and any spacelike hypersurface Σ that slices the support of T^{ab} . Then $(J^{ab})|_p = \int_{\Sigma} \overset{p}{\chi} [{}^a T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def} = \int_{\Sigma} \overset{o}{\chi} [{}^a T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def} + \int_{\Sigma} (\overset{p}{\chi} [{}^a - \overset{o}{\chi} [{}^a] T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def}$, where in the last step we have added and subtracted $\int_{\Sigma} \overset{o}{\chi} [{}^a T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def}$, which is a vector that we can understand to be defined at p . Notice that $(\overset{p}{\chi} [{}^a - \overset{o}{\chi} [{}^a])$ is a constant vector field: at any point q , it is just the vector “from p to q ” minus the vector “from o to q ”. Thus the field $(\overset{p}{\chi} [{}^a - \overset{o}{\chi} [{}^a])$ is given by the constant vector “from p to o ” at every point. This could be characterized as $(\overset{p}{\chi} [{}^a])|_o$ parallel transported to every point or alternatively as $-(\overset{o}{\chi} [{}^a])|_p$ parallel transported to every point. (See Fig. 2.) For clarity, we will again use the notation $(v^a)_{\parallel p}$ to represent the (global) vector field found by parallel transporting $(v^a)|_p$ to all points. In this notation, we have $(J^{ab})|_p = \int_{\Sigma} \overset{o}{\chi} [{}^a T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def} - \int_{\Sigma} (\overset{o}{\chi} [{}^a]_{\parallel p} T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def}$.




Figure 2: At any point q , $(\overset{p}{\chi} [{}^a - \overset{o}{\chi} [{}^a])|_q$ is the vector found by parallel transporting $-(\overset{o}{\chi} [{}^a])|_p$ to q .

Since $(\overset{o}{\chi} [{}^a])_{\parallel p}$ is a constant vector field, we can pull it out of the integral to write, $(J^{ab})|_p = \int_{\Sigma} \overset{o}{\chi} [{}^a T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def} - \left((\overset{o}{\chi} [{}^a]_{\parallel p} \int_{\Sigma} T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def} \right)|_p$. But $\left((\overset{o}{\chi} [{}^a]_{\parallel p} \right)|_p = (\overset{o}{\chi} [{}^a])|_p$ and $\int_{\Sigma} T^{bc} t_c \overset{\Sigma}{\epsilon}_{def} = P^b$, so we have $(J^{ab})|_p = \int_{\Sigma} \overset{o}{\chi} [{}^a T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def} - (\overset{o}{\chi} [{}^a P^b])|_p$. Moreover, in the present notation, $\int_{\Sigma} \overset{o}{\chi} [{}^a T^{b]c} t_c \overset{\Sigma}{\epsilon}_{def} = (J^{ab})|_o$. This means we can write $(J^{ab})|_p = \left((J^{ab})|_o - \overset{o}{\chi} [{}^a P^b] \right)|_p$. But p was arbitrary, so J^{ab} can be characterized in general as $J^{ab} = (J^{ab})|_o - \overset{o}{\chi} [{}^a P^b]$. Taking the action of ∇_a on both sides of this final expression yields $\nabla_a J^{bc} = -\delta_a^{[b} P^{c]}$. \square

3.5 Center of Mass

Now suppose additionally that (M, ∇) is geodesically complete.¹⁸ We can use the concepts already defined to describe the center of mass of T^{ab} .

Definition 3.10 *A set $A \subseteq M$ is spatially convex if and only if for all $p, q \in A$ for which there is a spacelike geodesic segment $\gamma : I \rightarrow M$ with endpoints p and q , $\gamma[I] \subseteq A$. For any tensor field $X_{b_1 \dots}$, let $X = \{\tilde{X} | \tilde{X} \text{ is spatially convex and } \text{supp}(X_{b_1 \dots}^{a_1 \dots}) \subseteq \tilde{X}\}$. Then the spatial convex hull¹⁹ of $X_{b_1 \dots}^{a_1 \dots}$, denoted $\text{ConvHull}(X_{b_1 \dots}^{a_1 \dots})$, is given by $\text{ConvHull}(X_{b_1 \dots}^{a_1 \dots}) = \bigcap X$.*

Proposition 3.11 *Let Σ be a spacelike hypersurface slicing the spatial convex hull of T^{ab} . There exists a unique point $q \in \Sigma$ such that $(J^{ab}t_b)|_q = \mathbf{0}$. Moreover, $q \in \text{ConvHull}(T^{ab})$.*

Proof. First we will prove that a point as described in the statement of the proposition exists. Fix some arbitrary $o \in \Sigma$ and consider $(J^{ab}t_b)|_o / (P^n t_n) = \int_{\Sigma} \overset{o}{\chi}{}^a T^{bc} t_b t_c \epsilon_{def}^{\Sigma} / (P^n t_n) = R^a$. Note that this expression is simply a definition of R^a —no claim has yet been made; moreover, $P^n t_n$ is just a scalar constant. We have used the fact that since $o \in \Sigma$, $\overset{o}{\chi}{}^a$ is spacelike on all of Σ to simplify this expression. R^a is a constant, spacelike vector field (spacelike because the integrand is spacelike over the entire domain of integration). We can then write $\int_{\Sigma} \overset{o}{\chi}{}^a T^{bc} t_b t_c \epsilon_{def}^{\Sigma} = R^a \int_{\Sigma} T^{bc} t_b t_c \epsilon_{def}^{\Sigma} = \int_{\Sigma} R^a T^{bc} t_b t_c \epsilon_{def}^{\Sigma}$ or $\int_{\Sigma} (\overset{o}{\chi}{}^a - R^a) T^{bc} t_b t_c \epsilon_{def}^{\Sigma} = \mathbf{0}$. But Σ is a spacelike hypersurface of a geodesically complete, simply connected classical spacetime, so it is a flat, three dimensional Euclidean manifold. Thus $\overset{o}{\chi}{}^a - R^a$ would be the position vector field centered at the point $q = o + R^a(o)$ (where we are using the natural affine structure of Euclidean space to represent points as a formal sum between a point and a vector, so a point p can be written as a sum of any point p' and a vector v from p' to p as $p = p' + v$),²⁰ if in fact there is such a point²¹ in Σ . But even if there is no such q in Σ , the vector field $\overset{o}{\chi}{}^a - R^a$ is well defined, and we can use the notation $\overset{o}{\chi}{}^a - R^a = \overset{q}{\chi}{}^a$ to describe a vector field on Σ without assuming that $q \in \Sigma$. Note, however, that if $q \in \Sigma$,

¹⁸As with simple connectedness, we will ultimately relax this condition by proceeding locally.

¹⁹At times, we will drop the ‘‘spatial,’’ but we will always mean the spatial convex hull.

²⁰For more on this notation, see Malament (2009).

²¹It is possible that Σ is bounded and $o + R^a(o)$ lies outside the bound of Σ , or that q has been excised from Σ .

then $(J^{ab}t_b)|_q = \mathbf{0}$ and q would be the desired point, so it only remains to show that $q \in \Sigma$ and we will have established existence.

We claim that there is such a point $q \in \Sigma$. To see why, first note that $\int_{\Sigma} \chi^a T^{bc} t_b t_c \epsilon_{def}^{\Sigma}$ is a positively weighted average of position vectors, and so it can only vanish if the position origin falls within the spacelike slice of the convex hull of T^{ab} over which the average is performed. (See, for instance, Benson (1966) for a proof of this well-known claim.) So $q \in \text{ConvHull}(T^{ab})$ (and *a fortiori*, $q \in M$, since M is geodesically complete). But Σ slices the spatial convex hull of T^{ab} , by hypothesis. So suppose there is no such q in Σ . Then we could define $\tilde{\Sigma} = \Sigma \cup \{q\}$. Since q is spacelike related to $o \in \Sigma$, $\tilde{\Sigma}$ is a spacelike hypersurface. Thus we have a spacelike hypersurface such that $\Sigma \subseteq \tilde{\Sigma}$ but $\Sigma \cap \text{ConvHull}(T^{ab}) \neq \tilde{\Sigma} \cap \text{ConvHull}(T^{ab})$, and so Σ does not slice $\text{ConvHull}(T^{ab})$, which is a contradiction. Thus, since $q \in \text{ConvHull}(T^{ab})$ and q is spacelike related to $o \in \Sigma$ (as it is by construction), $q \in \Sigma$.

It remains to show that q is unique. Suppose there were two such points, q and q' , where $q \neq q'$. Then $\int_{\Sigma} \chi^a T^{bc} t_b t_c \epsilon_{def}^{\Sigma} = \int_{\Sigma} \chi'^a T^{bc} t_b t_c \epsilon_{def}^{\Sigma} = \mathbf{0} = \int_{\Sigma} (\chi^a - \chi'^a) T^{bc} t_b t_c \epsilon_{def}^{\Sigma}$. Let R^a be as defined above and furthermore take Q^a be the unique constant vector field such that $q' = o + Q^a(o)$. Then we have $\int_{\Sigma} (\chi^a - \chi'^a) T^{bc} t_b t_c \epsilon_{def}^{\Sigma} = \int_{\Sigma} (R^a - Q^a) T^{bc} t_b t_c \epsilon_{def}^{\Sigma} = (R^a - Q^a) \int_{\Sigma} T^{bc} t_b t_c \epsilon_{def}^{\Sigma} = \mathbf{0}$. But $T^{bc} t_b t_c$ is nonvanishing and never negative by assumption (the first follows because T^{ab} is nonvanishing and the second by the mass condition), and so $\int_{\Sigma} T^{bc} t_b t_c \epsilon_{def}^{\Sigma} \neq 0$. Thus $R^a - Q^a = \mathbf{0}$ and $q = q'$. It follows that q is unique. \square

Prop. 3.11 allows us to speak of a single center of mass at a given time.

Definition 3.12 *Given a spacelike hypersurface Σ slicing the spatial convex hull of T^{ab} , we will call the unique $q \in \Sigma$ for which $(J^{ab}t_b)|_q = \mathbf{0}$ the center of mass of T^{ab} in Σ .*

Note finally that since $q \in \text{ConvHull}(T^{ab})$, we have a sense in which the center of mass is *inside* the worldtube of T^{ab} .

4 A Newtonian geodesic principle

We can now consider the motion of a particle in geometrized Newtonian theory. First, we require several lemmas.

Lemma 4.1 *Let (M, t_a, h^{ab}, ∇) be a classical spacetime, and suppose that M is oriented and simply connected and that (M, ∇) is geodesically complete. Assume that ∇ is flat. Let T^{ab} be a smooth symmetric tensor field on M satisfying: (1) the mass condition, (2) the conservation condition, and (3) given any spacelike hypersurface $\Sigma \subset M$, $\text{supp}(T^{ab}) \cap \Sigma$ is bounded. Let $G \subset M$ be the collection of center of mass points of T^{ab} . Then there is a smooth curve $(\gamma : I \rightarrow M) \in \Gamma$ (recall that Γ is the set of maximal integral curves of V^a) such that $G = \gamma[I]$.*

Proof. Consider any $g \in G$ and let $\gamma : I \rightarrow M$ be one of the (unique up to reparameterization) maximal integral curves of V^a passing through g . For concreteness, we can fix γ by supposing that $\gamma(0) = g$. By definition, $(J^{ab}t_b)|_g = \mathbf{0}$. Moreover, $V^n \nabla_n (J^{ab}t_b) = V^n (\delta_n^a P^b - \delta_n^b P^a) t_b = (V^a P^b - V^b P^a) t_b = \mathbf{0}$, so $J^{ab}t_b$ is constant along γ . Thus $J^{ab}t_b$ must vanish along all of γ and $\gamma[I] \subseteq G$. But $G \subseteq \gamma[I]$. Suppose otherwise. Then there would be some point $g' \in G$ such that $g' \notin \gamma[I]$. By the definition of a center of mass point, there must be some spacelike hypersurface Σ for which $g' \in \Sigma$. Since M is geodesically complete, there must be a spacelike hypersurface $\tilde{\Sigma}$ such that $\Sigma \subseteq \tilde{\Sigma}$ and $\gamma[I] \cap \tilde{\Sigma} \neq \emptyset$. But if $\gamma[I] \cap \tilde{\Sigma} \neq \emptyset$, there must be exactly one point in $p \in \gamma[I] \cap \tilde{\Sigma}$ because γ is a timelike curve. So $p \in G$ (since we already showed that $\gamma[I] \subseteq G$). By Prop. 3.11, there is exactly one center of mass point in any spacelike hypersurface slicing the convex hull of T^{ab} , so $p = g'$ and $g' \in \gamma[I]$, which is a contradiction. Thus $\gamma[I] = G$. \square

It follows immediately that in flat, simply connected, geodesically complete classical spacetimes, the path traced out by the center of mass of T^{ab} can always be reparameterized as a geodesic (so long as T^{ab} is conserved). In other words, Lemma 4.1 gives us a statement of Newton's first law, as a consequence of the mass condition, the conservation condition, and a condition on the boundedness of the body represented by T^{ab} . The second lemma is more complicated and involves a general classical spacetime.

Lemma 4.2 *Let (M, t_a, h^{ab}, ∇) be a classical spacetime and suppose M is simply connected. Moreover, suppose that $R^{abcd} = \mathbf{0}$ and $R^{ab}{}_{cd} = \mathbf{0}$.²² Let $\gamma : I \rightarrow M$ be a smooth timelike curve. Then there exists a flat derivative operator on M , $\overset{f1}{\nabla}$, that (1) is compatible with h^{ab} and t_a and (2) agrees with ∇ on γ .*

Proof. There are many flat derivative operators compatible with h^{ab} and t_a (Cf. Malament, 2010, Prop. 4.2.5).²³ Our strategy will be to construct one such operator (call it $\overset{f1}{\nabla}$) as in the proof to Prop. 4.2.5 and then keep it fixed as a reference. We will then construct a second operator that additionally satisfies (2) by making use of $\overset{f1}{\nabla}$.

Since $R^{abcd} = \mathbf{0}$ and $R^{ab}{}_{cd} = \mathbf{0}$, there exists (globally, since M is simply connected) a timelike vector field η^a that is rigid and non-rotating (i.e. $\nabla^a \xi^b = \mathbf{0}$). Let \hat{h}_{ab} be the spatial projection field relative to η^a (see Prop. 4.1.2) and define $\phi^a = \eta^n \nabla_n \eta^a$ and $\kappa_{ab} = \hat{h}_{n[b} \nabla_{a]} \eta^n$. We will take the reference derivative operator to be given by $\overset{f1}{\nabla} = (\nabla, \overset{01}{C}{}^a{}_{bc})$ where $\overset{01}{C}{}^a{}_{bc} = 2h^{am} t_{(b} \kappa_{c)m}$. As is shown in the proof of Prop. 4.2.5, this choice of derivative operator is flat and compatible with t_a and h^{ab} .

Prop. 4.2.5 shows that a second derivative operator/vector field pair $(\overset{f2}{\nabla}, \overset{2}{\phi}{}^a)$ will also be flat and compatible with h^{ab} and t_a iff $\nabla^a (\overset{2}{\phi}{}^b - \overset{1}{\phi}{}^b) = 0$ and $\overset{f2}{\nabla} = (\overset{f1}{\nabla}, \overset{12}{C}{}^a{}_{bc})$ where $\overset{12}{C}{}^a{}_{bc} = t_b t_c (\overset{2}{\phi}{}^a - \overset{1}{\phi}{}^a)$. Moreover, by Prop. 1.7.3, there must exist a symmetric tensor field $\overset{02}{C}{}^a{}_{bc}$ such that $\overset{f2}{\nabla} = (\overset{f1}{\nabla}, \overset{02}{C}{}^a{}_{bc})$. Indeed, $\overset{02}{C}{}^a{}_{bc} = \overset{01}{C}{}^a{}_{bc} + \overset{12}{C}{}^a{}_{bc}$.

One can write the required relation between $\overset{1}{\phi}{}^a$ and $\overset{2}{\phi}{}^a$ as $\overset{2}{\phi}{}^a = \overset{1}{\phi}{}^a + \psi^a$ where ψ^a is a covariant spacelike vector field satisfying $\nabla^b \psi^a = 0$. The condition that two derivative operators agree at a point p can be stated by demanding that the $\overset{02}{C}{}^a{}_{bc}$ field relating them vanishes at that point. Thus $\overset{f2}{\nabla}$ agrees with $\overset{f1}{\nabla}$ on γ just in case $\overset{02}{C}{}^a{}_{bc}$ vanishes on γ . This condition in turn holds just in case $\overset{01}{C}{}^a{}_{bc} + \overset{12}{C}{}^a{}_{bc} = 2h^{am} t_{(b} \kappa_{c)m} + t_b t_c \psi^a = 0$ on γ . Since η^a is timelike, $2t_{(b} \kappa_{c)}^a + t_b t_c \psi^a = 0$ on γ just in case $\eta^b \eta^c (2t_{(b} \kappa_{c)}^a + t_b t_c \psi^a) = 0$ on γ . (That this

²²What should one make of these conditions? Spatial flatness holds in any classical spacetime satisfying the geometrized version of Poisson's equation. The second condition, $R^{ab}{}_{cd} = \mathbf{0}$, is precisely the curvature condition necessary to recover standard Newtonian gravitation from geometrized Newtonian gravitation (Trautman, 1965; Malament, 2010). See section 2. This condition is strictly necessary for the argument given here to proceed.

²³All of the propositions of the form X.X.X cited in this proof are references to Malament (2010); we will refer to the proposition numbers directly and suppress further citations where no ambiguity can arise.

condition is necessary and sufficient follows from the fact that $\kappa^{ab} = 0$,²⁴ which implies that $h^{rb}(2t_{(b}\kappa_c)^a + t_b t_c \psi^a) = h^{rc}(2t_{(b}\kappa_c)^a + t_b t_c \psi^a) = 0$. Hence by the discussion following Prop. 4.1.1, if $\eta^b \eta^c (2t_{(b}\kappa_c)^a + t_b t_c \psi^a) = 0$ at p for *any* η^a , then $2t_{(b}\kappa_c)^a + t_b t_c \psi^a = 0$ identically at p .) But $\eta^b t_b = \eta^c t_c = 1$ and, as shown in the proof of Prop. 4.2.5, $2\kappa_a{}^b \eta^a = \overset{1}{\phi}{}^b$. Thus $\eta^b \eta^c (2t_{(b}\kappa_c)^a + t_b t_c \psi^a) = \overset{1}{\phi}{}^a + \psi^a$, and so $\overset{f2}{\nabla}$ agrees with ∇ on γ whenever $\psi^a = -\overset{1}{\phi}{}^a$ on γ . Note that this condition is equivalent to saying that, again on γ , $\overset{2}{\phi}{}^a = 0$.

As stated above, it is also necessary that $\nabla^b \psi^a = 0$ obtain. So we have two conditions on ψ^a (that it is constant in spacelike directions, and that it is the opposite of $\overset{1}{\phi}{}^a$ on γ). We claim that there is a field that meets both conditions. For any spacelike hypersurface Σ slicing the spatial convex hull of T^{ab} , let ψ^a be the vector field one finds by parallel transporting (relative to ∇) the vector $-\overset{1}{\phi}{}^a$ at the point where γ intersects Σ to all other points of Σ (this construction cannot produce ambiguities because we have assumed spatial flatness, and thus parallel transport in space is always path-independent, at least in a simply connected manifold). Then ψ^a is smooth, because $\overset{1}{\phi}{}^a$ is, and moreover, it satisfies both requirements. Thus $\overset{f2}{\nabla} = (\overset{f1}{\nabla}, t_b t_c \psi^a) = (\nabla, 2h^{am} t_{(b}\kappa_c)_m + t_b t_c \psi^a)$ is the required derivative operator. \square

It is important to note that the argument just given only permits one to find a flat derivative operator that agrees with ∇ on *timelike* curves. The argument fails for curves that intersect the same spacelike hypersurface more than once because ψ^a has to satisfy $\nabla^a \psi^b$. Once ψ^a has been chosen so that $\overset{f2}{\nabla}$ agrees with ∇ at *one* point on a given spacelike hypersurface, its value is fixed for the whole hypersurface, and so it cannot generally be made to agree with ∇ at any other point on the same spacelike hypersurface. This will complicate the proof of the result in the present paper, relative to the Geroch-Jang theorem, but it is not fatal, in large part because of the following result.

Lemma 4.3 *Let (M, t_a, h^{ab}, ∇) be an arbitrary classical spacetime, and suppose that M is*

²⁴Here is where the condition $R^{ab}{}_{cd} = \mathbf{0}$ is necessary. In a spatially flat classical spacetime, $R^{ab}{}_{cd} = \mathbf{0}$ iff there exists (at least locally) a rigid *and* non-rotating timelike field. If $R^{ab}{}_{cd} \neq \mathbf{0}$, then it is possible to find a rigid ($\nabla^a \eta^b = \mathbf{0}$) field, but not a non-rotating one; using this construction with a rigid but rotating field leads to $\kappa^{ab} = \nabla^{[a} \eta^{b]} \neq \mathbf{0}$, which means that in general one cannot find a flat derivative operator that agrees with ∇ on γ , at least by the argument offered here.

oriented and simply connected. Suppose also that $R^{abcd} = \mathbf{0}$. Let T^{ab} be a smooth symmetric tensor field on M satisfying: (1) the mass condition, (2) the conservation condition, and (3) given any spacelike hypersurface $\Sigma \subset M$, $\text{supp}(T^{ab}) \cap \Sigma$ is bounded. Suppose that Σ_1 and Σ_2 are spacelike hypersurfaces slicing the support of T^{ab} . Finally, let $\overset{f}{\nabla}$ be any flat derivative operator on M that is compatible with the spatial and temporal metrics. Then $t_a P^a(\Sigma_1) = t_a P^a(\Sigma_2)$, where $P^a(\Sigma_i)$ is defined relative to $\overset{f}{\nabla}$.

Proof. This result follows the proof of Prop. 3.2 closely. The most important thing to note is that here we assume that $\nabla_a T^{ab} = \mathbf{0}$, but not that $\overset{f}{\nabla}_a T^{ab} = \mathbf{0}$. Thus the argument that the integrand $\overset{f}{\nabla}_{[n} T^{ab} t_{|a} \overset{S}{\epsilon}_{b|cde]}$ vanishes fails. However, we now are considering a special case wherein $\kappa_a = t_a$. Without loss of generality, we can always choose to integrate relative to a set of basis fields in which t_a is a basis element. Then, by the Stokes' theorem argument given in the proof of Prop. 3.2, we have $t_a(P^a(\Sigma_1) - P^a(\Sigma_2)) = \int_S \overset{f}{\nabla}_{[n} T^{ab} t_{|a} \overset{S}{\epsilon}_{b|cde]}$. But $\overset{f}{\nabla}_{[n} T^{ab} t_{|a} \overset{S}{\epsilon}_{b|cde]}$ is an exterior derivative, and so it is invariant under different choices of covariant derivative operator. That is, we can write $\overset{f}{\nabla}_{[n} T^{ab} t_{|a} \overset{S}{\epsilon}_{b|cde]} = d_n(T^{ab} t_a \overset{S}{\epsilon}_{bcde}) = \nabla_{[n} T^{ab} t_{|a} \overset{S}{\epsilon}_{b|cde]}$, where in the last expression we are using the general curved derivative operator associated with the spacetime—relative to which T^{ab} is conserved. Again by reasoning present in the proof to Prop. 3.2, it can be shown that $\nabla_{[n} T^{ab} t_{|a} \overset{S}{\epsilon}_{b|cde]} = \nabla_b(T^{ab} t_a) \overset{S}{\epsilon}_{bcde}$. Since t_a is compatible with ∇ , we have $\nabla_b(T^{ab} t_a) = 0$. Thus $t_a(P^a(\Sigma_1) - P^a(\Sigma_2)) = 0$, or for any spacelike hypersurfaces slicing the support of T^{ab} , Σ_1 and Σ_2 , $P^a(\Sigma_1)t_a = P^a(\Sigma_2)t_a$. \square

It is now possible to state the general theorem concerning the Newtonian geodesic principle.²⁵

Theorem 4.4 *Let (M, t_a, h^{ab}, ∇) be a classical spacetime, and suppose that M is oriented and simply connected. Suppose also that $R^{abcd} = \mathbf{0}$ and $R^{ab}_{cd} = \mathbf{0}$. Let $\gamma : I \rightarrow M$ be a smooth imbedded curve. Suppose that given any open subset O of M containing $\gamma[I]$, there exists a smooth symmetric field $T^{ab} \in \mathfrak{T}^\bullet(M)$ with the following properties.*

1. T^{ab} satisfies the mass condition, i.e. whenever $T^{ab} \neq \mathbf{0}$, $T^{ab} t_a t_b > 0$;
2. T^{ab} satisfies the conservation condition, i.e. $\nabla_a T^{ab} = \mathbf{0}$;

²⁵The present proof is heavily indebted to the approximation scheme used in the proof of the Geroch-Jang theorem, though the framework in which their proof is presented is not immediately available in the classical context.

3. $\text{supp}(T^{ab}) \subset O$; and

4. there is at least one point in O at which $T^{ab} \neq \mathbf{0}$.

Then γ is a timelike curve that can be reparametrized as a geodesic.

Proof. We will consider three cases.

Case 1: First, suppose that γ is (everywhere) timelike. Let O be an open subset of M containing $\gamma[I]$ and let T^{ab} be a field meeting the requirements of the statement of the theorem. Since M is always locally geodesically complete, we can freely choose O so that there always exist geodesically complete spacelike hypersurfaces slicing the support of T^{ab} . By Lemma 4.2, there exists a flat derivative operator on M , $\overset{f}{\nabla}$, that is consistent with t_a and h^{ab} , and which agrees with ∇ on γ . For each spacelike hypersurface slicing the support of T^{ab} , Σ , it is possible to define $P^a(\Sigma)$ and $J^{ab}(\Sigma)$ (again, we can limit attention to geodesically complete hypersurfaces if necessary). These fields are defined relative to $\overset{f}{\nabla}$ in the sense that the parallel transport necessary to make sense of such integrals is performed relative to $\overset{f}{\nabla}$. Note that $P^a(\Sigma)$ and $J^{ab}(\Sigma)$ are globally defined fields; however, since T^{ab} is not necessarily conserved relative to $\overset{f}{\nabla}$, Props. 3.2 and 3.7 no longer hold and the fields are dependent on the choice of Σ . However, since each Σ is geodesically complete, Prop. 3.11 still holds for each Σ ; likewise Lemma 4.1, continues to hold for each of the $P^a(\Sigma)$ and $J^{ab}(\Sigma)$ fields individually (at least within a neighborhood of the unique center of mass point associated with Σ), relative to $\overset{f}{\nabla}$. Thus for each Σ , there is a geodesic $\overset{\Sigma}{\gamma}$ (relative to $\overset{f}{\nabla}$) that passes through the spatial convex hull of T^{ab} (relative to $\overset{f}{\nabla}$). One can think of these geodesics as the “unperturbed” paths of a particle: i.e. the paths a particle would take if the scalar field associated with $\overset{f}{\nabla}$ (the gravitational potential relative to $\overset{f}{\nabla}$) suddenly vanished at the time associated with a given Σ .

As has already been mentioned, T^{ab} is not necessarily conserved relative to $\overset{f}{\nabla}$. However, $\overset{f}{\nabla}_a T^{ab} = (\overset{f}{\nabla}_a - \nabla_a)T^{ab}$ is given by a smooth field (a sum of products of the $\overset{02}{C}{}^a{}_{bc}$ field constructed in Lemma 4.2 and T^{ab}) that vanishes on γ , since by construction the two operators agree there. Thus, for any constant scalar field $\alpha > 0$, one can make $|\overset{f}{\nabla}_a T^{ab} t_b| < \alpha$

everywhere by shrinking the support of T^{ab} (which is always possible because a suitable T^{ab} exists for *any* neighborhood of γ).

Let Σ_1 and Σ_2 be any two appropriate spacelike hypersurface slicing the support of T^{ab} and consider the fields $J^{ab}(\Sigma_1)t_a$ and $J^{ab}(\Sigma_2)t_b$. The curves $\overset{\Sigma_1}{\gamma}$ and $\overset{\Sigma_2}{\gamma}$ consist of the points at which $J^{ab}(\Sigma_1)t_a$ and $J^{ab}(\Sigma_2)t_a$ vanish, respectively. Now let Σ be some other appropriate spacelike hypersurface slicing the support of T^{ab} , and let $p \in \Sigma$. The field $J^{ab}(\Sigma_1)t_a$ (for instance) at p can be interpreted as the vector pointing from p to o , where o is the point at which $\overset{\Sigma_1}{\gamma}$ intersects Σ . Note that this interpretation makes sense because (1) Σ is always a flat space with Euclidean affine structure and (2) $J^{ab}t_a$ is always spacelike (as can be seen immediately by the symmetry properties of J^{ab}). This means that at any p in an appropriate Σ , the vector $(J^{ab}(\Sigma_1) - J^{ab}(\Sigma_2))t_a$ represents the vector from p to o , minus the vector from p to o' (where o' is the point at which $\overset{\Sigma_2}{\gamma}$ intersects Σ), which is just the vector from o' to o . Note that this difference is independent of p , but dependent on the spacelike hypersurface containing p . So we can define a (spacelike) vector field $\eta^a = (J^{ab}(\Sigma_1) - J^{ab}(\Sigma_2))t_b$ whose spatial length at any point p in a spacelike hypersurface slicing the support of T^{ab} represents the distance between the points at which $\overset{\Sigma_1}{\gamma}$ and $\overset{\Sigma_2}{\gamma}$ intersect that spacelike hypersurface.

Our goal will be to show that the spatial length of η^a can be made arbitrarily small everywhere. To see this, note that since η^a is always spacelike, there exists a vector β_a such that $\eta_a = h^{ab}\beta_b$. The spatial length of η^a is then given by $(h^{ab}\beta_a\beta_b)^{1/2}$. Pick an arbitrary point $p \in M$ and consider $h^{ab}\beta_a\beta_b = \beta_a\eta^a$ at p . By definition of the terms involved, this last expression can be written in terms of a constant basis $\overset{1}{\sigma}_a, \dots, \overset{4}{\sigma}_a$ (relative to $\overset{f}{\nabla}$), so that

$$h^{ab}\beta_a\beta_b = \sum_{i=1}^4 \overset{i}{\beta} \left(\int_{\Sigma_1} \overset{p}{\chi}^{[aT^b]c} \overset{i}{\sigma}_a t_b t_c \overset{\Sigma_1}{\epsilon}_{def} - \int_{\Sigma_2} \overset{p}{\chi}^{[aT^b]c} \overset{i}{\sigma}_a t_b t_c \overset{\Sigma_2}{\epsilon}_{def} \right). \quad (4.1)$$

By the Stokes' theorem reasoning in the proof of Prop. 3.2, we can construct a submanifold S with Σ_1 and Σ_2 forming partial boundaries, such that,

$$h^{ab}\beta_a\beta_b = \sum_{i=1}^4 \beta^i \int_S \overset{f}{\nabla}_{[n} \overset{p}{\chi}^{[a} T^{b]c} \overset{i}{\sigma}_{|a} t_b \epsilon_{c|def]}^S. \quad (4.2)$$

Again by the reasoning of the proof of Prop. 3.2, we can show that $\overset{f}{\nabla}_c(\overset{p}{\chi}^{[a} T^{b]c} \overset{i}{\sigma}_a t_b) = \overset{p}{\chi}^{[a}(\overset{f}{\nabla}_c T^{b]c}) \overset{i}{\sigma}_a t_b$. This final expression, meanwhile, represents a scalar field that can be made as small as one likes by shrinking the support of T^{ab} . It follows that the righthand side of Eq. (4.2) can be made arbitrarily small. And so, for any positive scalar field α , one can choose O so that $h^{ab}\beta_a\beta_b < \alpha$.

It follows that for any two appropriate spacelike hypersurfaces Σ_1 and Σ_2 , the geodesics $\overset{\Sigma_1}{\gamma}$ and $\overset{\Sigma_2}{\gamma}$ can be made arbitrarily close to one another in the sense that, given any two appropriate spacelike hypersurfaces slicing the support of T^{ab} , Σ_1 and Σ_2 , and any open set A containing $\overset{\Sigma_1}{\gamma}[I]$, we can choose T^{ab} so that $\overset{\Sigma_2}{\gamma}[I] \subset A$ as well. Moreover, for each Σ , $\overset{\Sigma}{\gamma}$ passes through the intersection of the spatial convex hull (relative to $\overset{f}{\nabla}$) of T^{ab} and Σ , and so we can conclude that the image of the original curve, $\gamma[I]$, is arbitrarily close to a geodesic (relative to $\overset{f}{\nabla}$), in the same sense. This last result is only possible if γ can itself be reparameterized as a geodesic (relative to $\overset{f}{\nabla}$). Finally, since $\overset{f}{\nabla}$ agrees with ∇ on γ , then γ must be a geodesic relative to ∇ as well, up to reparameterization.

Case 2: Now suppose γ is (everywhere) spacelike. We claim that there exist open sets containing $\gamma[I]$ for which there does not exist a smooth symmetric field $T^{ab} \in \mathfrak{T}^\bullet(M)$ satisfying conditions 1-4. Suppose that for any open set containing $\gamma[I]$, such a field did exist. We know that there always exists a flat derivative operator on M , so let $\overset{f}{\nabla}$ be any such flat derivative operator. Since γ is everywhere spacelike, there must be some spacelike hypersurface Σ such that $\gamma[I] \subseteq \Sigma$.

First, suppose that Σ can be chosen to be bounded. Then we can also freely choose a neighborhood O of γ which is also bounded. Since M is simply connected, it admits a

global time function, $t : M \rightarrow \mathbb{R}$, which is unique up to an additive constant. We can choose O so that there is some value t' of the time function with the following property: if Σ' is a spacelike hypersurface whose time value is t' , Σ' satisfies $\Sigma' \cap O = \emptyset$. It follows that T^{ab} vanishes on Σ' , and thus that $P^a(\Sigma') = \mathbf{0}$ (where the integrals are performed relative to the arbitrary flat derivative operator $\overset{f}{\nabla}$). Thus $P^a(\Sigma')t_a = 0$. Meanwhile, by the mass condition, we know that $P^a(\Sigma)t_a > 0$. Now we can use a slightly modified²⁶ version of the argument of Lemma 4.3. Since O is bounded, we can freely choose some third (timelike) hypersurface Σ'' (adjusting our choices of O and Σ if necessary) s.t. $\Sigma'' \cap O = \emptyset$, and such that $\Sigma \cup \Sigma' \cup \Sigma''$ forms the boundary of a four dimensional submanifold of M , S (where we reverse the orientation of, say, Σ' so that S is outwardly oriented). We can thus apply the Stokes' theorem argument given in the proofs of Prop. 3.2 and Lemma 4.3 to show that $P^a(\Sigma)t_a = P^a(\Sigma')t_a$, which is a contradiction.

Now suppose that Σ cannot be chosen to be bounded. For simplicity, we will assume that Σ can be chosen so that it extends to spatial infinity in all directions.²⁷ Choose O so that it has the following property: in the limit of spatial infinity, the “temporal height” of O goes to zero (see Fig. 3). Here's one way (of many) to make this idea precise. Without loss of generality, choose the time function t so that for any $s \in I$, $t(\gamma(s)) = 0$. Let ϖ be any (fixed) timelike geodesic passing through Σ . Then given any point p in a spacelike hypersurface intersecting ϖ , we can define a distance function $d : M \rightarrow \mathbb{R}$ relative to ϖ as the (spatial) distance from ϖ to p . We can then define an open set $O = \{p \in M \mid |t(p)| < a \text{ and } |d(p)t(p)| < 1\}$, for some constant real number a chosen so that ϖ intersects all of the simultaneity slices of M with time values from $-a$ to a . Note first that $\Sigma \subset O$, so $\gamma[I] \subset O$. Moreover, for any $p \in O - \Sigma$, there exists a spacelike hypersurface Σ' for which $p \in \Sigma'$ and Σ' slices O (since the restriction of O to any spacelike hypersurface except Σ is bounded by

²⁶Modified because by our definition, Σ' does not slice the support of T^{ab} , since $\Sigma' \cap \text{supp}(T^{ab}) = \emptyset$. But in this special case the argument still goes through.

²⁷We are ignoring the case where Σ is unbounded, but not necessarily in all directions. The argument given here is intended to be representative: it can be extended to include these more complicated cases by, for instance, choosing O so that the temporal height of its closure would vanish at any boundary of Σ .




Figure 3: An example in three dimensions of an open set O whose “temporal height” goes to zero at spatial infinity, and which contains a spacelike hypersurface. In the case where γ is restricted to a single spacelike hypersurface, Σ , the depicted set can be chosen to contain γ . Thus there must be some point $p \in O$ at which T^{ab} is non-vanishing. By smoothness, we can assume $p \notin \Sigma$. One can then choose a cylinder (as shown) whose bottom (or top) slices O , but which doesn’t intersect the set elsewhere. The argument in Lemma 4.3 then yields a contradiction. (See Case 2 in the text.)

construction).

From here the argument is similar to the bounded case. For any given a , there exist spacelike hypersurfaces Σ_{\pm} such that for any $p \in \Sigma_+$, $t(p) > a$, and for any $p \in \Sigma_-$, $t(p) < -a$. These are necessarily such that $\Sigma_{\pm} \cap O = \emptyset$. It follows that T^{ab} vanishes on Σ_{\pm} , and thus that $P^a(\Sigma_{\pm}) = \mathbf{0}$ (where the integrals are performed relative to the arbitrary flat derivative operator $\overset{f}{\nabla}$). Thus $P^a(\Sigma_{\pm})t_a = 0$. Meanwhile, we know there must be some point $p \in O$ at which $T^{ab} \neq \mathbf{0}$. We can freely suppose that $t(p) \neq 0$ (because if $t(p) = 0$, there necessarily exists a neighborhood around p in which $T^{ab} \neq \mathbf{0}$, since T^{ab} is smooth, and which must include points whose time values are greater and less than 0). Suppose without loss of generality that $t(p) > 0$ (if $t(p) < 0$, simply reverse the temporal order of the ensuing argument—we have already chosen O so that there are temporally prior, non-intersecting spacelike hypersurfaces). Since $p \in O - \Sigma$, we know there's a spacelike hypersurface Σ' that contains p and slices O . By the mass condition and the smoothness of T^{ab} , we know that $P^a(\Sigma')t_a > 0$. Now we can use Stokes' theorem as immediately above by connecting Σ' and Σ_+ to reason to a contradiction. Thus γ cannot be spacelike.

Case 3: So far we have shown that if γ is everywhere timelike then it must be (reparametrizable as) a geodesic, and that γ cannot be everywhere spacelike. The final case concerns curves that are sometimes timelike and sometimes spacelike. Given case 1, it is sufficient to show that if γ satisfies the assumptions of the theorem and is timelike at at least one point, then it is timelike everywhere. Suppose it isn't—i.e., suppose there is at least one point q at which γ is spacelike. Let $s_1 \in I$ be such that γ is timelike at $\gamma(s_1)$ and let $s_2 \in I$ be such that γ is spacelike at $\gamma(s_2)$. Let ξ^a be the tangent field to γ . We can define a scalar field on γ by $\alpha = \xi^a t_a$. α can be understood as a smooth function $\alpha : I \rightarrow \mathbb{R}$ defined by $\alpha(s) = \alpha \circ \gamma(s) = (\xi^a t_a)|_{\gamma(s)}$. Since γ is timelike at $\gamma(s_1)$, we know that $\alpha(s_1) > 0$; likewise, since γ is spacelike at $\gamma(s_2)$, $\alpha(s_2) = 0$. Since α is just a smooth function on the reals, however, we know that there must be a number $t \in I$ such that $\alpha(t) > 0$, but for which $(\frac{d}{ds}\alpha)(t) \neq 0$. But by definition of ξ^a , $\frac{d}{ds}\alpha(s) = (\xi^a|_{\gamma(s)})(\alpha) = \xi^a \nabla_a \alpha = t_b \xi^a \nabla_a \xi^b$. So at $\gamma(t)$,

we know that $(t_b \xi^a \nabla_a \xi^b)|_{\gamma(t)} \neq 0$, and that $\xi^a t_a > 0$.

So γ is timelike at $\gamma(t)$, which means (since γ is smooth and imbedded) that there must be an open neighborhood Q of $\gamma(t)$ such that the restriction of $\gamma[I]$ to Q is timelike. (Why? Since γ is smooth, there must be an open neighborhood $J \subseteq I$ of t such that $\gamma[J]$ is timelike. And since γ is imbedded, there must be an open subset Q of M such that $\gamma[J] = \gamma[I] \cap Q$. So the restriction of $\gamma[I]$ to Q is timelike and contains $\gamma(t)$.) We can freely choose Q so that it is simply connected. Note that since γ is such that for any neighborhood of γ , there exists a smooth symmetric field T^{ab} satisfying conditions 1-4, it follows that for any *sub*-neighborhood Q' of Q containing $\gamma[I] \cap Q$, there also exists a smooth symmetric field T^{ab} such that the restriction of T^{ab} to Q satisfies conditions 1-4, relative to Q' . (Why? Extend Q' to a neighborhood O of all of γ in any way at all, so long as $O \cap Q = Q'$. Then a field T^{ab} satisfying conditions 1-4 relative to O is guaranteed to exist by the assumptions of the theorem; the restriction of T^{ab} to Q automatically inherits conditions 1-3. And by the conservation of mass argument given in Lemma 4.3, if T^{ab} is non-vanishing anywhere within O , as it must be, then it is possible to show by a series of flux integrals that it is non-vanishing along the length of the curve, and so T^{ab} must be non-vanishing somewhere in Q' .) But then if we take Q as a submanifold of M and take the restriction of γ to Q as a timelike curve, case 1 applies and γ must be a geodesic everywhere in Q . It follows that at $\gamma(t) \in Q'$, $(\xi^a \nabla_a \xi^b)|_{\gamma(t)} = 0$, which is a contradiction (since we showed that $(t_b \xi^a \nabla_a \xi^b)|_{\gamma(t)} \neq 0$). And so γ must be timelike everywhere. \square

Mathematically, theorem 4.4 differs from the Geroch-Jang theorem in at least two important ways.²⁸ First, it requires two curvatures conditions: spatial flatness and $R^{ab}_{cd} = \mathbf{0}$. Spatial flatness follows immediately from the geometrized version of Poisson's equation; however, the Geroch-Jang theorem does not require one to assume Einstein's equation (or any curvature conditions following from it), and so the requirement of spatial flatness is perhaps

²⁸As mentioned in footnote 6, we are deferring questions concerning interpretational differences between the two theorems in the contexts of their respective spacetime theories to a later paper. Here we are only concerned with the mathematical structures of the two theorems.

a defect of Theorem 4.4. The second curvature condition, meanwhile, is necessary to recover standard Newtonian gravitation from the geometrized theory. Without it, it is possible to find a more general “Newtonian” theory (see Künzle (1976); Ehlers (1981); Malament (2010)), but with a vector potential replacing the scalar potential of standard Newtonian gravitation, and with a universal rotation field affecting the behavior of this vector potential. Strictly speaking, this generalized Newtonian theory is the classical limit of GR. However, the present argument for Lemma 4.2 fails if we relax this second curvature condition, and so we are not guaranteed that a flat derivative operator exists that agrees with the curved operator on γ . It is quite likely (we believe) that a different argument can be given to show that such a derivative operator does exist, in which case it would be possible to relax the condition $R^{ab}{}_{cd} = \mathbf{0}$ in Theorem 4.4. We would like to note, however, that insofar as we were interested in the status of the geodesic principle in *Newtonian* physics (rather than in some generalized Newtonian physics), $R^{ab}{}_{cd} = \mathbf{0}$ is a perfectly reasonable requirement: it holds just in case a geometrized Newtonian spacetime admits a standard Newtonian representation. It is part of what makes a classical spacetime Newtonian.

Secondly, Theorem 4.4 requires the assumption that the underlying manifold M be simply connected; the Geroch-Jang theorem, however, does not seem to require any such global topological assumptions.²⁹ The reason that simple-connectedness is required here is that vector integration in a classical spacetime, at least as we have developed it, requires simple connectedness to ensure a unique result for the integral (since otherwise, parallel transport is not necessarily globally unique). Geroch and Jang use Killing fields to avoid this problem entirely; however, one does not have access to timelike Killing fields, even locally, in a classical spacetime. However, there are two simple corollaries available that extend the result to a more general case. The first uses the fact that *any* manifold is *locally* simply connected; the second uses the fact that a geodesic need only be locally extremal.

²⁹It is possible that there *is* a global topological assumption lurking in the background of the Geroch-Jang theorem, though we do not see where it enters. Reconstructing the proof of the Geroch-Jang theorem in detail is beyond the scope of this paper.

Corollary 4.5 *Let (M, t_a, h^{ab}, ∇) be a classical spacetime, and suppose that M is oriented. Suppose also that $R^{abcd} = \mathbf{0}$ and $R^{ab}_{cd} = \mathbf{0}$. For any $p \in M$, there exists a neighborhood of p , Q , such that if (1) $\gamma : I \rightarrow Q$ is a smooth curve, and (2) for any open subset O of Q containing $\gamma[I]$ there exists a smooth symmetric field $T^{ab} \in \mathfrak{T}^\bullet(M)$ satisfying conditions 1-4 of Theorem 4.4, then γ is a timelike curve that can be reparametrized as a geodesic (segment).*

Corollary 4.6 *Let (M, t_a, h^{ab}, ∇) be a classical spacetime, and suppose that M is oriented. Suppose also that $R^{abcd} = \mathbf{0}$ and $R^{ab}_{cd} = \mathbf{0}$. Let $\gamma : I \rightarrow M$ be a smooth curve with the following property: for any $p \in \gamma[I]$, there exists a neighborhood of p , Q , such that in any open subset O of Q containing the restriction of $\gamma[I]$ to Q there exists a smooth symmetric field $T^{ab} \in \mathfrak{T}^\bullet(M)$ satisfying conditions 1-4 of Theorem 4.4, then γ is a timelike curve that can be reparametrized as a geodesic.*

Corollary 4.5 precisifies a sense in which *local* geodesic motion has the status of a general theorem in geometrized Newtonian gravitation. Corollary 4.6, meanwhile, shows that there is a sense in which one can relax the requirement of simple connectedness and still find a global result. It is important to emphasize, however, that we do not get something for nothing: the additional requirement that conditions 1-4 hold in the neighborhood of any point of the curve is a substantial strengthening of the conditions required for Theorem 4.4. In particular, if one assumes simply that conditions 1-4 hold in any open set O containing the curve mentioned in Theorem 4.4, then it need not follow that T^{ab} is non-vanishing in any open set containing only *part* of the curve, as would be necessary to generalize Corollary 4.6 further.

References

- Benson, Russell V. (1966). *Euclidean geometry and convexity*. New York: McGraw-Hill.
- Boothby, William M. (2003). *An Introduction to Differentiable Manifolds and Riemannian Geometry*. 2nd edition. San Diego: Academic Press.
- Cartan, Elie (1923). “Sur les variétés à connexion affine, et la théorie de la relativité généralisée (première partie).” *Annales scientifiques de l’École Normale Supérieure*, 40, 325–412.
- Cartan, Elie (1924). “Sur les variétés à connexion affine, et la théorie de la relativité généralisée (première partie) (suite).” *Annales scientifiques de l’École Normale Supérieure*, 41, 1–25.

- Dixon, W. G. (1964). “A Covariant Multipole Formalism for Extended Test Bodies in General Relativity.” *Il Nuovo Cimento*, 34(2), 317–339.
- Duval, Ch. and H. P. Künzle (1978). “Dynamics of Continua and Particles from General Covariance of Newtonian Gravitation Theory.” *Reports on Mathematical Physics*, 13(3).
- Ehlers, Jürgen (1981). “Über den Newtonschen Grenzwert der Einsteinschen Gravitationstheorie.” *Grundlagen Probleme der Modernen Physik*, Ed. Jürgen Nitsch, Joachim Pfarr, and Ernst-Walther Stachow. Zurich: Bibliographisches Institut.
- Ehlers, Jürgen and Robert Geroch (2004). “Equation of motion of small bodies in relativity.” *Annals of Physics*, 309, 232–236.
- Einstein, A., L. Infeld, and B. Hoffman (1938). “The Gravitational Equations and the Problem of Motion.” *Annals of Mathematics*, 39(1), 65–100.
- Friedrichs, Kurt O. (1927). “Eine Invariante Formulierung des Newtonschen Gravitationsgesetzes und der Grenzüberganges vom Einsteinschen zum Newtonschen Gesetz.” *Mathematische Annalen*, 98, 566–575.
- Geroch, Robert and Pong Soo Jang (1975). “Motion of a body in general relativity.” *Journal of Mathematical Physics*, 16(1), 65.
- Künzle, H. P. (1976). “Covariant Newtonian Limit of Lorentz Space-Times.” *General Relativity and Gravitation*, 7(5), 445–457.
- Malament, David (1986). “Newtonian Gravity, Limits, and the Geometry of Space.” *From Quarks to Quasars*, Ed. Robert Colodny. Pittsburgh: University of Pittsburgh Press.
- Malament, David B. (2009). “Notes on Geometry and Spacetime.”
- Malament, David B. (2010). “Relativistic and Newtonian Spacetime Structure.”
- Taub, A. H. (1962). “On Thomas’ Result Concerning the Geodesic Hypothesis.” *Proceedings of the National Academy of the USA*, 48(9), 1570–1571.
- Thomas, T. Y. (1962). “On the Geodesic Hypothesis in the Theory of Gravitation.” *Proceedings of the National Academy of the USA*, 48(9), 1567–1569.
- Trautman, Andrzej (1965). “Foundations and Current Problem of General Relativity.” *Lectures on General Relativity*, Ed. Stanley Deser and Kenneth W. Ford. Englewood Cliffs, NJ: Prentice-Hall.
- Suijlekom, Walter D. van and Eli J. Hawkins (2009). “Crash course in differential geometry.”